

R1321T**Polyclonal Antibody to Goat IgG F(ab')₂ -TRITC-**

Alternate names:	Goat Immunoglobulin G
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Host:	Rabbit
Immunogen:	Goat IgG F(ab') ₂ fragment.
Format:	State: Lyophilized purified Ig fraction. Purification: Immunoaffinity chromatography. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer. Label: TRITC – Tetramethylrhodamine isothiocyanate ; Molecular Weight 444 daltons) <i>Absorption / Emission:</i> 550 nm / 570 nm <i>Molar Ratio:</i> 2.8 moles TRITC per mole of Rabbit IgG. Reconstitution: Restore with 1.0 ml of deionized water (or equivalent).
Applications:	Suitable for Immunomicroscopy (1/500-1/2,500) and Flow cytometry or FACS analysis (1/2,000-1/10,000) as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG, Goat IgG F(ab') ₂ and Goat Serum. No reaction was observed against Goat IgG F(c) or Human Serum Proteins.
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. Shelf life: One year from despatch.
General Readings:	J.A. Titus, et al. J. Immunol. Methods 50; 193, 1982. (Conjugation)