

**R1387P****Fab Fragments of Donkey anti-Goat IgG [H&L]**

<b>Alternate names:</b>	Goat Immunoglobulin G
<b>Quantity:</b>	1 mg
<b>Concentration:</b>	1.0 mg/ml (by UV absorbance at 280 nm)
<b>Host:</b>	Donkey
<b>Immunogen:</b>	Goat IgG whole molecule.
<b>Format:</b>	<b>State:</b> Liquid (sterile filtered) Fab fragments. <b>Purification:</b> Immunoaffinity chromatography. <b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 0.01% (w/v) Sodium Azide as preservative.
<b>Applications:</b>	Suitable for Immunoprecipitation, Immunodiffusion, conjugation and most immunological methods requiring lot-to-lot consistency, high titer and specificity. Recommended dilutions for ELISA: 1:10,000-1:50,000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Donkey Serum. No reaction was observed against anti-Pepsin or anti-Donkey IgG F(c).
<b>Storage:</b>	Store the antibody (undiluted) at 2-8°C for one month or (in aliquots) at -20°C for longer. Dilute only prior to immediate use. Avoid cycles of freezing and thawing. Shelf life: One year from despatch.