

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046 Web: www.ebioscience.com E-mail: info@ebioscience.com

Product Information

Contents: Fluorescein isothiocyanate (FITC) anti-human CD209

(DC-SIGN)


Catalog Number: 11-2099 Sizes: 25 tests, 100 tests

Formulation: Phosphate buffer pH 7.2, 150 mM NaCl, 0.09% NaN₃, 0.2% BSA Storage Conditions: Store at 4°C.

DO NOT FREEZE.

LIGHT-SENSITIVE MATERIAL.

Clone: eB-h209 I sotype: Rat IgG2a, κ HLDA No.: N/A



Human monocyte-derived immature dendritic cells (culture with GM-CSF and IL-4) were stained with FITC-eB-h209 (shaded histogram). Rat IgG2a isotype control is demonstrated by the open histogram. Cells were run on a FACSCalibur and large cells were gated.

| Available Formats of This Product | | | | |
|-----------------------------------|--|-------------|--------------|-----------------------|
| Cat. No. | Format | Excite (nm) | Emit (nm) | Reported Applications |
| 11-2099 | FITC anti-human CD209 (DC-SIGN) | 488 | 518 | FC |
| 12-2099 | PE anti-human CD209 (DC-SIGN) | 488 | 575 | FC |
| 13-2099 | Biotin anti-human CD209 (DC-SIGN) | N/A | N/A | FC |
| 14-2099 | Affinity Purified anti-human CD209 (DC-SIGN) | N/A | N/A | FC IP |
| 17-2099 | APC anti-human CD209 (DC-SIGN) | 633 | 660 | FC |

Description

The eB-h209 monoclonal antibody reacts with human CD209, also known as DC-SIGN, a 44 kDa type II transmembrane protein. DC-SIGN contains a C-type lectin binding domain and binds ICAM-3, ICAM-2, and HIV virus. Human dendritic cells preferentially express DC-SIGN. It has been postulated that DC-SIGN serves as a receptor for capture, trafficking, and transmission of HIV to T cells and supports primary immune response. eB-h209 was developed against a C-terminal peptide of human DC-SIGN.

Usage

For research use only, not for diagnostic or therapeutic use. The eB-h209 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The eB-h209 antibody has been pre-titrated and tested by flow cytometric analysis of cultured human dendritic cells and peripheral blood leukocytes. This can be used at 20 µl per 100 µl blood (or per 1 million cells in 100 µl total staining volume).

Related Products

Cat. 12-2099 PE anti-human CD209 (DC-SIGN) (clone eB-h209)
Cat. 13-2099 Biotin anti-human CD209 (DC-SIGN) (clone eB-h209)

Cat. 14-2099 Affinity Purified anti-human CD209 (DC-SIGN) (clone eB-h209)

Cat. 17-2099 APC anti-human CD209 (DC-SIGN) (clone eB-h209)

Cat. 11-4321 FITC Rat IgG2a Isotype Control

References

eBioscience (2001). Unpublished results.

Geijtenbeek, T.B, D.S. Douglas, et al. (2000) "DC-SIGN, a Dendritic Cell-Specific HIV-1-Binding protein that Enhances trans-Infection of T cells." <u>Cell</u> 100(5): 587-597.

Geijtenbeek, T.B, R Torensma, et al. (2000). "Identification of DC-SIGN, a Novel Dendritic Cell-Specific ICAM-3 Receptor that Supports Primary Immune Responses." Cell 100(5): 575-585.

Geijtenbeek, T.B, D.J. Krooshop, et al. (2000). "DC-SIGN-ICAM-2 Interaction Mediates Dendritic Cell Trafficking." Nat. Immunol. 1 (4):353-357

Pohlmann, S, F Baribaud, et al. (2001). "DC-SIGN Interactions with Human Immunodeficiency Virus type 1 and 2 and Simian Immunodeficiency Virus." J Virol. 75(10):4664-4672

 $\label{eq:convergence} \text{Copyright} \ @ \ 2000\text{-}2005 \ \text{eBioscience, Inc.}$ Product For Research Use Only: Not for further distribution without written consent.