

Product Information

Contents: Fluorescein isothiocyanate (FITC) anti-mouse FcεRIα (FcεRI-α, high affinity IgE receptor)

Catalog Number: 11-5898

Sizes: 50 µg, 100 µg, 500 µg

Formulation: Phosphate buffer pH 7.2,
150 mM NaCl, 0.09% NaN₃


Storage Conditions: Store at 4°C.

DO NOT FREEZE.

LIGHT-SENSITIVE MATERIAL.

Clone: MAR-1

Isotype: Armenian Hamster IgG



Staining of MC/9 cells with 0.06 µg of FITC Armenian hamster IgG isotype control (cat. 11-4444) (open histogram) or 0.06 µg of FITC MAR-1 (colored histogram). Total viable cells were used for analysis.

Available Formats of This Product

Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
11-5898	FITC anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor)	488	518	FC
12-5898	PE anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor)	488	575	FC
13-5898	Biotin anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor)	N/A	N/A	FC
14-5898	Affinity Purified anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor)	N/A	N/A	FA FC IH/F IHC/P IP WB
16-5898	Functional Grade* Purified anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor)	N/A	N/A	FA FC

*Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/µg.
Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

Description

The MAR-1 monoclonal antibody reacts with the FcεRIα subunit, an IgE-binding subunit lacking signal-transducing ability. FcεRIα is expressed on mast and basophil cells and is up-regulated by the presence of IgE. FcεRIα forms a tetrameric complex with one β and two γ subunits. The β and γ subunits possess immunoreceptor tyrosine-based activation motifs (ITAM). The FcεRI complex plays an important role in triggering IgE-mediated allergic reactions.

Usage

For research use only, not for diagnostic or therapeutic use. The MAR-1 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The MAR-1 antibody has been tested by flow cytometric analysis of the MC/9 cell line (a mouse mast cell line). This can be used at less than or equal to 0.125 µg per million cells in a 100 µl total staining volume. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Related Products

Cat. 11-4444 FITC Armenian Hamster IgG Isotype Control (clone n/a)

Cat. 12-5898 PE anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor) (clone MAR-1)
Cat. 13-5898 Biotin anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor) (clone MAR-1)
Cat. 14-5898 Affinity Purified anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor) (clone MAR-1)
Cat. 16-5898 Functional Grade Purified anti-mouse FcεRI alpha (FcεRIα, FcεRI-a, FcεRI-alpha, high affinity IgE receptor) (clone MAR-1)

References

Personal Correspondence

Yamaguchi M., K. Hirai, A. Komiya, M. Miyamasu, Y. Furumoto, R. Teshima, K. Ohta, Y. Morita, S. J. Galli, C. Ra, K. Yamamoto. (2001) Regulation of Mouse Mast Cell Surface FcεRI expression by dexamethasone. *Int Immunol* 13(7):843-51.

Copyright © 2000-2005 eBioscience, Inc.
Product For Research Use Only: Not for further distribution without written consent.

