

Product Information

Contents: Phycoerythrin (PE) anti-mouse CD8a (Ly-2)

Catalog Number: 12-0081

Sizes: 50 ug, 100 ug, 200 ug, 500 ug

Formulation: Phosphate buffer pH 7.2,
150 mM NaCl, 0.09% NaN₃


Storage Conditions: Store at 4°C.

DO NOT FREEZE.

LIGHT-SENSITIVE MATERIAL.

Clone: 53-6.7

Isotype: Rat IgG2a, κ



Staining of mouse thymocytes (left) and splenocytes (right) with FITC anti-CD4 (GK1.5) (cat.11-0041) and 0.125 μg PE anti-mouse CD8 (53-6.7). Total viable cells were used for analysis.

Available Formats of This Product

Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
10-0081	APC-Cy7 Rat anti-mouse CD8 (alpha subunit; CD8a)	633	760	FC
11-0081	FITC Rat anti-mouse CD8 (alpha subunit; CD8a)	488	518	FC
12-0081	PE Rat anti-mouse CD8 (alpha subunit; CD8a)	488	575	FC
13-0081	Biotin Rat anti-mouse CD8 (alpha subunit; CD8a)	N/A	N/A	FC
14-0081	Affinity Purified Rat anti-mouse CD8 (alpha subunit; CD8a)	N/A	N/A	FA FC IH/F IP
15-0081	PE-Cy5 Rat anti-mouse CD8 (alpha subunit; CD8a)	488	670	FC
16-0081	Functional Grade* Purified Rat anti-mouse CD8 (alpha subunit; CD8a)	N/A	N/A	FA FC
17-0081	APC Rat anti-mouse CD8 (alpha subunit; CD8a)	633	660	FC
19-0081	Cy5 Rat anti-mouse CD8 (alpha subunit; CD8a)	633	670	FC
25-0081	PE-Cy7 Rat anti-mouse CD8 (alpha subunit; CD8a)	488	760	FC
30-0081	DISCONTINUED - Allophycocyanin-Cy5.5 (APC-Cy5.5) anti-mouse CD8a (Ly-2)	633	690	FC

*Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/μg.
Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

Description

The 53-6.7 monoclonal antibody reacts with the mouse CD8a molecule. CD8a is an approximately 32-34 kDa cell surface receptor expressed either as a heterodimer with the CD8 β chain (CD8 αβ) or as a homodimer (CD8 αα). A majority of thymocytes and a subpopulation of mature αβ TCR T cells express CD8 αβ while γδ TCR T cells, a subpopulation of intestinal intraepithelial lymphocytes (IELs) and dendritic cells express CD8 αα. CD8 binds to MHC class I and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells.

Usage

For research use only, not for diagnostic or therapeutic use. The 53-6.7 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The 53-6.7 antibody has been tested by flow cytometric analysis of mouse thymocyte and splenocyte suspensions. This can be used at less than or equal to 0.25 μg per million cells in a 100 μl total staining volume. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Related Products

Cat. 10-0081 APC-Cy7 Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 11-0081 FITC Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 13-0081 Biotin Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 14-0081 Affinity Purified Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 15-0081 PE-Cy5 Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 16-0081 Functional Grade Purified Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 17-0081 APC Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 19-0081 Cy5 Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 25-0081 PE-Cy7 Rat anti-mouse CD8 (alpha subunit; CD8a) (clone 53-6.7)
Cat. 12-4321 PE Rat IgG2a Isotype Control

References

Ledbetter, J. A. and L. A. Herzenberg (1979). "Xenogeneic monoclonal antibodies to mouse lymphoid differentiation antigens." Immunol Rev 47: 63-90.

Ledbetter, J. A., R. V. Rouse, et al. (1980). "T cell subsets defined by expression of Lyt-1,2,3 and Thy-1 antigens. Two-parameter immunofluorescence and cytotoxicity analysis with monoclonal antibodies modifies current views." J Exp Med 152(2): 280-95.

Copyright © 2000-2005 eBioscience, Inc.
Product For Research Use Only: Not for further distribution without written consent.

