

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046 Web: www.ebioscience.com

Web: www.ebioscience.com E-mail: info@ebioscience.com

#### **Product Information**

Contents: Phycoerythrin (PE) anti-human CD49d (Integrin a<sub>4</sub>)

Catalog Number: 12-0499 Sizes: 25 tests, 100 tests


Formulation: Phosphate buffer pH 7.2, 150 mM NaCl, 0.09% NaN<sub>3</sub>, 0.2% BSA Storage Conditions: Store at 4°C.

DO NOT FREEZE.

LIGHT-SENSITIVE MATERIAL.

Clone: 9F10

Isotype: Mouse IgG1, κ HLDA No.: V S215



Staining of normal human peripheral blood cells with PE mouse IgG1 isotype control (cat. 12-4719) (open histogram) or PE 9F10 (colored histogram). Cells in the lymphocyte gate were used for analysis.

Available Formats of This Product				
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
12-0499	PE anti-human CD49d (Integrin alpha 4)	488	575	FC
14-0499	Affinity Purified anti-human CD49d (Integrin alpha 4)	N/A	N/A	FA FC IHC
16-0499	Functional Grade* Purified anti-human CD49d (Integrin alpha 4)	N/A	N/A	FA FC
*	L CradaTM (ECTM). Azida-fron starila-filtared and andatovin < 0.001 n	~ /···~		

<sup>\*</sup>Functional Grade<sup> $\mathbb{M}$ </sup> (FG $\mathbb{M}$ ): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/ $\mathbb{M}$ 9. Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

## Description

The 9F10 monoclonal antibody reacts with human CD49d, the 150 kDa integrin  $a_4$  subunit. The complex of CD49d non-covalently associated with integrin  $\beta_1$  (CD29), also known as VLA-4, is a receptor for fibronectin and VCAM-1 (CD106). This complex is expressed by thymocytes, peripheral lymphocytes, monocytes and eosinophils. CD49d also associates with integrin  $\beta_7$  and binds to the Mucosal Addressin Cell-Adhesion Molecule-1 (MadCAM-1).

## Usage

For research use only, not for diagnostic or therapeutic use. The 9F10 antibody has been reported for use in flow cytometric analysis.

# **Applications Tested**

The 9F10 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 20 µl per 100 µl blood (or per 1 million cells in 100 µl total staining volume).

## Related Products

Cat. 14-0499 Affinity Purified anti-human CD49d (Integrin alpha 4) (clone 9F10)

Cat. 16-0499 Functional Grade Purified anti-human CD49d (Integrin alpha 4) (clone 9F10)

Cat. 12-4714 Phycoerythrin (PE) Mouse IgG1, K Isotype Control

#### References

Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York.

Copyright © 2000-2005 eBioscience, Inc.
Product For Research Use Only: Not for further distribution without written consent.