

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046 Web: www.ebioscience.com

Web: www.ebioscience.com E-mail: info@ebioscience.com

Product Information

Contents: Phycoerythrin (PE) anti-mouse CD103 (Integrin a_{IFI}

chain)

Catalog Number: 12-1031 Sizes: 50 ug, 100 ug, 200 ug

Formulation: Phosphate buffer pH 7.2,

150 mM NaCl, 0.09% NaN₃


Storage Conditions: Store at 4°C.


DO NOT FREEZE.

LIGHT-SENSITIVE MATERIAL.

Clone: 2E7

Isotype: Armenian Hamster IgG





Staining of C57BI/6 lymph node cell suspension with FITC anti-CD3e (cat.11-0031) with staining buffer (autofluorescence) (left) or 05\ microg of PE 2E7 (right). Total viable cells were used for analysis.

Available Formats of This Product					
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications	
11-1031	FITC anti-mouse CD103 (Integrin alpha-IEL chain)	488	518	FC	
12-1031	PE anti-mouse CD103 (Integrin alpha-IEL chain)	488	575	FC	
13-1031	Biotin anti-mouse CD103 (Integrin alpha-IEL chain)	N/A	N/A	FC	
14-1031	Affinity Purified anti-mouse CD103 (Integrin alpha-IEL chain)	N/A	N/A	FA FC IH/F IP	
16-1031	Functional Grade* Purified anti-mouse CD103 (Integrin alpha-IEL chain)	N/A	N/A	FA FC	

^{*}Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/µg. Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

Description

The 2E7 monoclonal antibody reacts with mouse CD103, the $a_{\rm E}$ integrin. CD103 non-covalently associates with integrin β_7 and is expressed on all intraepithelial lymphocytes, a small subset of peripheral lymphocytes, and dendritic epidermal T cells (DEC). Epithelial cell antigen E-cadherin binds to CD103 and mediates homing of lymphocytes to the intestinal epithelium.

Usage

For research use only, not for diagnostic or therapeutic use. The 2E7 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The 2E7 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to $0.5 \mu g$ per million cells in a $100 \mu l$ total staining volume. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Related Products

Cat. 11-1031	FITC anti-mouse CD103 (Integrin alpha-IEL chain) (clone 2E7)
Cat. 13-1031	Biotin anti-mouse CD103 (Integrin alpha-IEL chain) (clone 2E7)
Cat. 14-1031	Affinity Purified anti-mouse CD103 (Integrin alpha-IEL chain) (clone 2E7)
Cat. 16-1031	Functional Grade Purified anti-mouse CD103 (Integrin alpha-IEL chain) (clone 2E7)
Cat. 12-4914	Phycoerythrin (PE) Golden Syrian Hamster IgG Isotype Control (clone n/a)

References

Lefrancois, L., T. A. Barrett, et al. (1994). "Developmental expression of the alpha IEL beta 7 integrin on T cell receptor gamma delta and T cell receptor alpha beta T cells." <u>Eur J Immunol</u> 24(3): 635-40.

Copyright © 2000-2005 eBioscience, Inc.
Product For Research Use Only: Not for further distribution without written consent.