


Product Information

Contents: Phycoerythrin (PE) anti-human interleukin-4 (IL-4)  
Catalog Number: 12-7048  
Sizes: 25 tests, 50 ug, 100 ug  
Formulation: Phosphate buffer pH 7.2,  
150 mM NaCl, 0.09% NaN<sub>3</sub>  
Storage Conditions: Store at 4°C.  
DO NOT FREEZE.  
LIGHT-SENSITIVE MATERIAL  
Clone: MP4-25D2  
Isotype: Rat IgG1, κ



Human peripheral blood mononuclear cells were stimulated with PMA and ionomycin in the presence of Brefeldin A for 5 hours. (Left) The cells were surface stained with PE anti-human CD3 (UCHT1) and intracellularly stained with FITC anti-human IL-4 (MP4-25D2). (Right) The cells were surface stained with FITC anti-human CD3 and intracellularly stained with PE anti-human IL-4.

Available Formats of This Product				
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
11-7048	FITC anti-human IL-4 (Interleukin-4)	488	518	IC
12-7048	PE anti-human IL-4 (Interleukin-4)	488	575	IC
13-7048	Biotin anti-human IL-4 (Interleukin-4)	N/A	N/A	ELISA det ELISPOT det
14-7048	Affinity Purified anti-human IL-4 (Interleukin-4)	N/A	N/A	IC
16-7048	Functional Grade* Purified anti-human IL-4 (Interleukin-4)	N/A	N/A	BA
17-7048	APC anti-human IL-4 (Interleukin-4)	633	660	IC

\*Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/µg.  
Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

Description

The MP4-25D2 antibody reacts with human interleukin-4 (IL-4), a 15-19 kDa cytokine secreted by Th2 cells.

Usage

For research use only, not for diagnostic or therapeutic use. MP4-25D2 has been reported for use in intracellular flow cytometric analysis.

Applications Tested

The PE conjugated MP4-25D2 is offered in 2 formats:

- µg size: has been tested by intracellular flow cytometric analysis and can be used at less than or equal to 0.5 µg/million cells (100 µl blood). It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.
- test size: has been pre-titrated and tested by intracellular flow cytometric analysis and can be used at 20 µl per million cells (100 µl blood). The pre-titrated test size contains 0.2% BSA.

Related Products

Cat. 12-4301 PE Rat IgG1 Isotype Control

## References

---

- Ramanathan, L., R. Ingram, et al. (1993). Immunochemical mapping of domains in human interleukin 4 recognized by neutralizing monoclonal antibodies. *Biochemistry* 32(14): 3549-56.
- Abrams, J. S., M. G. Roncarolo, et al. (1992). Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol Rev* 127: 5-24.

Copyright © 2000-2005 eBioscience, Inc.  
Product For Research Use Only: Not for further distribution without written consent.

