

Product Information

Contents: Phycoerythrin (PE) anti-human CD95 Ligand (Fas Ligand, CD178)

Catalog Number: 12-9919

Sizes: 25 tests, 100 tests

Formulation: Phosphate buffer pH 7.2, 150 mM NaCl, 0.09% NaN₃, 0.2% BSA

Storage Conditions: Store at 4°C.


DO NOT FREEZE.

LIGHT-SENSITIVE MATERIAL.

Clone: NOK-1

Isotype: Mouse IgG1, κ

HLDA No.: N/A



Surface staining of human FasL transfected cells with anti-human CD178 (NOK-1) PE. Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.

Available Formats of This Product

Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
12-9919	PE anti-human CD95 Ligand (Fas Ligand, CD178)	488	575	FC
13-9919	Biotin anti-human CD95 Ligand (Fas Ligand, CD178)	N/A	N/A	FC
14-9919	Affinity Purified anti-human CD95 Ligand (Fas Ligand, CD178)	N/A	N/A	FA FC IP
16-9919	Functional Grade* Purified anti-human CD95 Ligand (Fas Ligand, CD178)	N/A	N/A	FA FC

*Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/μg.
Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

Description

The NOK-1 monoclonal antibody reacts with human Fas (CD95) Ligand, a 40 kDa type II transmembrane glycoprotein. FasL is a member of the TNF family and is expressed by neutrophils, monocytes, and activated T cells and NK cells. The interaction of FasL with its receptor (CD95, Fas) induces Fas-mediated killing of lymphocytes. Human FasL is cleaved from the surface by matrix metalloproteinases (MMPs), resulting in a 26 kDa soluble form. Therefore for optimal detection of surface FasL on activated peripheral blood cells, incubation of cells with an MMP inhibitor is recommended.

Usage

For research use only, not for diagnostic or therapeutic use. NOK-1 has been reported for use in flow cytometric analysis.

Applications Tested

The NOK-1 antibody has been pre-titrated and tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at 20 μl per 100 μl blood (or per 1 million cells in 100 μl total staining volume).

Related Products

- Cat. 12-4714 Phycoerythrin (PE) Mouse IgG1, K Isotype Control
- Cat. 13-9919 Biotin anti-human CD95 Ligand (Fas Ligand, CD178) (clone NOK-1)
- Cat. 14-9919 Affinity Purified anti-human CD95 Ligand (Fas Ligand, CD178) (clone NOK-1)
- Cat. 16-9919 Functional Grade Purified anti-human CD95 Ligand (Fas Ligand, CD178) (clone NOK-1)

References

Suda, T., H. Hashimoto, et al. (1997). "Membrane Fas ligand kills human peripheral blood T lymphocytes, and soluble Fas ligand

blocks the killing." J Exp Med 186(12): 2045-50.

Kayagaki, N., A. Kawasaki, et al. (1995). "Metalloproteinase-mediated release of human Fas ligand." J Exp Med 182(6): 1777-83.

Tanaka, M., T. Suda, et al. (1995). "Expression of the functional soluble form of human fas ligand in activated lymphocytes." Embo J 14(6): 1129-35.

Copyright © 2000-2005 eBioscience, Inc.

Product For Research Use Only: Not for further distribution without written consent.

