


Product Information

Contents: Biotin anti-mouse IgD
Catalog Number: 13-5993
Sizes: 50 ug, 100 ug, 500 ug
Formulation: Phosphate buffer pH 7.2,
150 mM NaCl, 0.1% NaN₃
Storage Conditions: Store at 4°C.
DO NOT FREEZE.
Clone: 11-26c (11-26)
Isotype: Rat IgG2a, κ



Surface staining of mouse splenocytes with anti-mouse IgD (11-26c) PE. Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.

Available Formats of This Product

Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
11-5993	FITC anti-mouse IgD	488	518	FC
12-5993	PE anti-mouse IgD	488	575	FC
13-5993	Biotin anti-mouse IgD	N/A	N/A	FC
14-5993	Affinity Purified anti-mouse IgD	N/A	N/A	FC

Description

The 11-26.c monoclonal antibody reacts with the δ heavy chain of mouse IgD. It does not react with other classes of mouse immunoglobulin including IgA, IgG or IgM. IgD is expressed by peripheral mature B cells. 11-26.c does not activate B cells.

Usage

For research use only, not for diagnostic or therapeutic use. The 11-26 antibody has been reported for use in flow cytometric analysis to detect the surface IgD on mouse B cells.

Applications Tested

The 11-26c (11-26) antibody has been tested by flow cytometric analysis of mouse splenocytes. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Related Products

Cat. 11-4317 Streptavidin-FITC (Fluorescein isothiocyanate)
Cat. 12-4317 Streptavidin-PE (Phycoerythrin)
Cat. 17-4317 Streptavidin Allophycocyanin (SA-APC)
Cat. 13-4321 Biotin Rat IgG2a Isotype Control
Cat. 24-5093 Polyclonal anti-mouse IgD antiserum (activating) (clone Polyclonal)
Cat. 11-5993 FITC anti-mouse IgD (clone 11-26c (11-26))
Cat. 12-5993 PE anti-mouse IgD (clone 11-26c (11-26))
Cat. 14-5993 Affinity Purified anti-mouse IgD (clone 11-26c (11-26))

References

Nitschke L, Kosco MH, Koehler G, and Lamers MC. 1993. Immunoglobulin D-deficient mice can mount normal immune responses to

thymus-independent and -dependent antigens. Proc Natl. Acad. Sci. USA. 90:1887-1891.

Ishihara K, Wood Jr. WJ, Wall R, Sakaguchi N, Michnoff C, Tucker PW, and Kincade PW. 1993. Multiple B29 containing complexes on murine B lymphocytes. Common and stage-restricted Ig-associated polypeptide chains. J. Immunol. 150: 2253 - 2262.

Copyright © 2000-2005 eBioscience, Inc.
Product For Research Use Only: Not for further distribution without written consent.

