

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046 Web: www.ebioscience.com E-mail: info@ebioscience.com

### **Product Information**

Contents: Affinity Purified anti-human CD11a (LFA-1a)

Catalog Number: 14-0119

Sizes: 25 ug, 100 ug


Formulation: Phosphate buffer pH 7.2,

150 mM NaCl, 0.09% NaN<sub>3</sub>

Storage Conditions: Store at 4°C. Avoid repeated freeze/thaw cycles.

Clone: HI111

I sotype: Mouse IgG1, κ HLDA No.: IV N231



Profile of purified mAb HI111 (CD11a) staining on human peripheral blood lymphocytes.

Available Formats of This Product				
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
11-0119	FITC anti-human CD11a	488	518	FC
12-0119	PE anti-human CD11a	488	575	FC
14-0119	Affinity Purified anti-human CD11a	N/A	N/A	FA FC IHC
16-0119	Functional Grade* Purified anti-human CD11a	N/A	N/A	FA FC

<sup>\*</sup>Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/μg. Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

# Description

The HI111 monoclonal antibody reacts with human CD11a, the 180 kDa integrin  $a_L$ , also known as the lymphocyte function associated antigen-1 (LFA-1) a chain. LFA-1, formed by non-covalent association of CD11a with CD18 (integrin  $\beta_2$ ), serves as an important adhesion molecule involved in lymphocyte and granulocyte function. CD54 (ICAM-1), CD102 (ICAM-2), and CD50 (ICAM-3) are ligands for LFA-1. CD11a is expressed by all leukocytes.

## Usage

For research use only, not for diagnostic or therapeutic use. The HI111 antibody has been reported for use in flow cytometric analysis, and immunohistochemical staining. It has also been reported in blocking of CD11a function in mixed lymphocyte reactions. (Please use Functional Grade purified HI111, cat. 16-0119, in functional assays.)

# Applications Tested

The HI111 antibody has been tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at less than or equal to 1 µg per 100 µl blood (or per 1 million cells in 100 µl total staining volume). It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

#### **Related Products**

Cat. 11-0119 FITC anti-human CD11a (clone HI111)

Cat. 12-0119 PE anti-human CD11a (clone HI111)

Cat. 16-0119 Functional Grade Purified anti-human CD11a (clone HI111)

Cat. 11-4011 FITC Anti-Mouse IgG

Cat. 13-4013 Biotin Anti-Mouse IgG (clone Polyclonal)

Cat. 11-4317 Streptavidin-FITC (Fluorescein isothiocyanate)

Cat. 12-4317 Streptavidin-PE (Phycoerythrin)

Cat. 17-4317 Streptavidin Allophycocyanin (SA-APC)

Cat. 14-4714 Affinity Purified Mouse IgG1, K Isotype Control

## References

Knapp, W., B. Dorken, et al. eds. 1989. Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press. New York.

 $\label{lem:copyright} \hbox{ $\mathbb{Q}$ 2000-2005 eBioscience, Inc.}$  Product For Research Use Only: Not for further distribution without written consent.