

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046 Web: www.ebioscience.com

Web: www.ebioscience.com E-mail: info@ebioscience.com

Product Information

Contents: Affinity Purified anti-mouse CD94

Catalog Number: 14-0941 Sizes: 50 ug, 100 ug, 500 ug


Formulation: Phosphate buffer pH 7.2,

150 mM NaCl, 0.09% NaN₃

Storage Conditions: Store at 4°C. Avoid repeated freeze/thaw cycles.

Clone: 18d3

Isotype: Rat IgG2a, κ



Surface staining of mouse splenocytes with anti-mouse CD94 (18d3) FITC (left), and PE (right). Appropriate isotype controls were used (open histogram). Total viable cells were used for analysis.

Available Formats of This Product				
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications
11-0941	FITC anti-mouse CD94	488	518	FC
12-0941	PE anti-mouse CD94	488	575	FC
13-0941	Biotin anti-mouse CD94	N/A	N/A	FC
14-0941	Affinity Purified anti-mouse CD94	N/A	N/A	FC

Description

The 18d3 monoclonal antibody reacts with mouse CD94, a 70 kDa type II transmembrane glycoprotein. CD94 belongs to the C-type lectin superfamily and is present as a heterodimer with NKG2 on the surface. CD94 is expressed by NK cells, a subset of T cells, and NKT cells and plays an important role in adhesion and activation of NK cell lineage. DBA/2J mice are naturally CD94-deficient and do not express cell-surface CD94/NKG2A receptors, even on neonatal NK cells.

Usage

For research use only, not for diagnostic or therapeutic use. The 18d3 antibody has been reported for use in flow cytometric analysis.

Applications Tested

The 18d3 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.25 µg per million cells in a 100 µl total staining volume. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Related Products

FITC anti-mouse CD94 (clone 18d3) Cat. 11-0941 Cat. 12-0941 PE anti-mouse CD94 (clone 18d3) Cat. 13-0941 Biotin anti-mouse CD94 (clone 18d3) Streptavidin-FITC (Fluorescein isothiocyanate) Cat. 11-4317 Cat. 12-4317 Streptavidin-PE (Phycoerythrin) Cat. 17-4317 Streptavidin Allophycocyanin (SA-APC) Cat. 14-4321 Affinity Purified Rat IgG2a Isotype Control Cat. 11-4811 FITC Anti-Rat IgG

Biotin Anti-Rat IgG (clone Polyclonal)

Cat. 13-4813

Vance, R. E., A. M. Jamieson, et al. (1999). "Recognition of the class Ib molecule Qa-1(b) by putative activating receptors CD94/NKG2C and CD94/NKG2E on mouse natural killer cells." J Exp Med 190(12): 1801-12.

Vance, R. E., J. R. Kraft, et al. (1998). "Mouse CD94/NKG2A is a natural killer cell receptor for the nonclassical major histocompatibility complex (MHC) class I molecule Qa-1(b)." J Exp Med 188(10): 1841-8.

Vance, R. E., D. M. Tanamachi, et al. (1997). "Cloning of a mouse homolog of CD94 extends the family of C-type lectins on murine natural killer cells." Eur J Immunol 27(12): 3236-41.

 ${\it Copyright} \ @ \ 2000-2005 \ eBioscience, \ Inc. \\ {\it Product For Research Use Only: Not for further distribution without written consent.} \\$