

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046

Web: www.ebioscience.com E-mail: info@ebioscience.com

Product Information

Contents: Functional Grade Purified anti-human CD3

Catalog Number: 16-0039

Sizes: 50 ug, 500 ug

Formulation: Phosphate buffer pH 7.2,

150 mM NaCl, No NaN₃


Storage Conditions: Store at 4°C. Avoid repeated freeze/thaw cycles.

KEEP CONTENT STERILE.

Endotoxin Level: Less than 0.001 ng/ug antibody, as determined

by the LAL assay. Clone: HIT3a

Isotype: Mouse IgG2a, к HLDA No.: V 5T-CD03.05



Surface staining of normal human peripheral blood cells with antihuman CD3 (HIT3a) PE. Autofluorescence is indicated by open histogram. Cells in the lymphocyte population were used for analysis.

Available Formats of This Product					
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications	
11-0039	FITC anti-human CD3	488	518	FC	
12-0039	PE anti-human CD3	488	575	FC	
14-0039	Affinity Purified anti-human CD3	N/A	N/A	FC	
16-0039	Functional Grade* Purified anti-human CD3	N/A	N/A	FA FC	

*Functional Grade™ (FG™): Azide-free, sterile-filtered, and endotoxin < 0.001 ng/µg. Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

Description

The HIT3a monoclonal antibody reacts with human CD3e, a 20 kDa subunit of the TCR complex. Along with the other CD3 subunits γ and δ , the ϵ chain is required for proper assembly, trafficking and surface expression of the TCR complex. CD3 is expressed by thymocytes in a developmentally regulated manner and by all mature T cells. Crosslinking of TCR with HIT3a initiates an intracellular biochemical pathway resulting in cellular activation and proliferation.

Usage

For research use only, not for diagnostic or therapeutic use. The HIT3a antibody has been reported for use in flow cytometric analysis.

Applications Tested

The HIT3a antibody has been tested by flow cytometric analysis of human peripheral blood leukocytes. This can be used at less than or equal to $0.25~\mu g$ per $100~\mu l$ blood (or per 1 million cells in $100~\mu l$ total staining volume). It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Related Products

Cat. 11-0037 FITC anti-human CD3 (clone OKT3) Cat. 12-0037 PE anti-human CD3 (clone OKT3)

Cat. 14-0037 Affinity Purified anti-human CD3 (clone OKT3)

Cat. 16-0037 Functional Grade Purified anti-human CD3 (clone OKT3)

Cat. 10-0038 APC-Cy7 anti-human CD3 (clone UCHT1)
Cat. 11-0038 FITC anti-human CD3 (clone UCHT1)
Cat. 12-0038 PE anti-human CD3 (clone UCHT1)

Cat. 13-0038 Biotin anti-human CD3 (clone UCHT1)

Cat. 14-0038	Affinity Purified anti-human CD3 (clone UCHT1)
Cat. 15-0038	PE-Cy5 anti-human CD3 (clone UCHT1)
Cat. 16-0038	Functional Grade Purified anti-human CD3 (clone UCHT1)
Cat. 17-0038	APC anti-human CD3 (clone UCHT1)
Cat. 25-0038	PE-Cy7 anti-human CD3 (clone UCHT1)
Cat. 11-0039	FITC anti-human CD3 (clone HIT3a)
Cat. 12-0039	PE anti-human CD3 (clone HIT3a)
Cat. 14-0039	Affinity Purified anti-human CD3 (clone HIT3a)
Cat. 11-4011	FITC Anti-Mouse IgG
Cat. 13-4013	Biotin Anti-Mouse IgG (clone Polyclonal)
Cat. 11-4317	Streptavidin-FITC (Fluorescein isothiocyanate)
Cat. 12-4317	Streptavidin-PE (Phycoerythrin)
Cat. 17-4317	Streptavidin Allophycocyanin (SA-APC)
Cat. 16-4724	Functional Grade Purified Mouse IgG2a, K Isotype Control

References

McMichael, A.J., P.C.L. Beverly, et al. eds. (1987). Leucocyte Typing III: White Cell Differentiation Antigens. Oxford University Press. New York.

Knapp, W., B. Dorken, et al. eds. (1989). Leucocyte Typing IV: White Cell Differentiation Antigens. Oxford University Press. New York.

Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York.

 $\label{localization} \text{Copyright} \ @ \ 2000\text{-}2005 \ \text{eBioscience, Inc.} \\ \text{Product For Research Use Only: Not for further distribution without written consent.} \\$