

Tel: 888.999.1371 or 858.642.2058

Fax: 858.642.2046 Web: www.ebioscience.com E-mail: info@ebioscience.com

#### **Product Information**

Contents: Functional Grade Purified anti-mouse CD106 (VCAM-1)

Catalog Number: 16-1061 Sizes: 50 ug, 100 ug, 500 ug

Formulation: Phosphate buffer pH 7.2,

150 mM NaCl, No NaN<sub>3</sub>

Storage Conditions: Store at 4°C. Avoid repeated freeze/thaw cycles.

KEEP CONTENT STERILE.

Endotoxin Level: Less than 0.001 ng/ug antibody, as determined by the LAL assay.

Clone: 429

Isotype: Rat IgG2a, κ

Available Formats of This Product					
Cat. No.	Format	Excite (nm)	Emit (nm)	Reported Applications	
11-1061	FITC anti-mouse CD106 (VCAM-1)	488	518	FC	
13-1061	Biotin anti-mouse CD106 (VCAM-1)	N/A	N/A	FC	
14-1061	Affinity Purified anti-mouse CD106 (VCAM-1)	N/A	N/A	FA FC IHC IP	
16-1061	Functional Grade* Purified anti-mouse CD106 (VCAM-1)	N/A	N/A	FA FC	
*Function	al Grade <sup>TM</sup> (FG <sup>TM</sup> ): Azide-free, sterile-filtered, and endotoxin $< 0$ .	.001 ng/µg.			

Purified: Contains azide, not sterile-filtered, and not endotoxin tested.

#### Description

The 429 monoclonal antibody reacts with mouse CD106 (Vascular Cell Adhesion Molecule-1, VCAM-1), a 110 kDa transmembrane glycoprotein expressed by myeloid lineage and bone marrow stromal cells. Endothelial cells constitutively express low levels of CD106 and upregulate it upon cytokine stimulation. CD106 binds to integrin  $a_4\beta_1$  (VLA-4, CD49d/CD29) and Integrin a4b7 (LPAM-1) and these interactions in the bone marrow and thymus are important for early lymphocyte and myeloid development. Cytokine-mediated upregulation of CD106 on endothelial cells suggests a role for this antigen in the inflammatory response.

# Usage

For research use only, not for diagnostic or therapeutic use. The 429 antibody has been reported for use in flow cytometric analysis. It has also been reported in blocking of CD106 in functional studies.

## **Applications Tested**

The 429 antibody has been tested by flow cytometric analysis of mouse bone marrow cell suspensions. This can be used at less than or equal to 0.25 µg per million cells in a 100 µl total staining volume. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

## **Related Products**

Cat. 11-1061	FITC anti-mouse CD106 (VCAM-1) (clone 429)
Cat. 13-1061	Biotin anti-mouse CD106 (VCAM-1) (clone 429)
Cat. 14-1061	Affinity Purified anti-mouse CD106 (VCAM-1) (clone 429)
Cat. 11-4317	Streptavidin-FITC (Fluorescein isothiocyanate)
Cat. 12-4317	Streptavidin-PE (Phycoerythrin)
Cat. 17-4317	Streptavidin Allophycocyanin (SA-APC)

Cat. 16-4321 Functional Grade Purified Rat IgG2a Isotype Control

Cat. 11-4811 FITC Anti-Rat IgG

Cat. 13-4813 Biotin Anti-Rat IgG (clone Polyclonal)

### References

Kinashi, T., Y. St. Pierre, et al. (1995). Expression of glycophosphatidylinositol-anchored and -non-anchored isoforms of vascular cell adhesion molecule 1 in murine stromal and endothelial cells. J Leukoc Biol 57(1): 168-73.

 $\label{lem:copyright} \hbox{ $\mathbb{Q}$ 2000-2005 eBioscience, Inc.}$  Product For Research Use Only: Not for further distribution without written consent.