

Caco-2 Lysate

Catalog No.: AL00022PU-S

Quantity: 0.1 mg

Concentration: 2mg/ml

Format: **State:** Liquid

Purity: Caco-2 lysate was prepared by homogenization in modified RIPA buffer (150 mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 ug/ml of aprotinin, 5 ug/ml of leupeptin.) Cell debris was removed by centrifugation. The Caco-2 lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% beta-mercaptoethanol.

Applications: Western Blot - 10µg to 20µg per lane are recommended for mini gels. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.

Description: CACO-2 cells total protein lysate

Storage: Store at 2-8°C for continuous use or (in aliquots) at -70°C for longer. Avoid repeated freezing and thawing. Shelf life: 6 month from despatch.