

AM00007BT-N

Monoclonal Antibody to Amyloid beta (free N-term, low APP reactive) - Biotin

Alternate names:	Amyloid beta peptide
Quantity:	0.1 mg
Concentration:	0.5 mg/ml
Background:	The beta-amyloid peptide (beta A4), proteolytically released from the amyloid precursor protein (APP), is the principal component of senile plaques in Alzheimer's disease. Cleavage of APP by alpha-secretase or alternatively by beta-secretase leads to generation and extracellular release of soluble APP peptides, S-APP-alpha and S-APP-beta, respectively, and the retention of corresponding membrane-anchored C-terminal fragments, C83 and C99. Subsequent processing of C83 by gamma-secretase yields P3 peptides. This is the major secretory pathway and is nonamyloidogenic. Alternatively, presenilin/nicastrin-mediated gamma-secretase processing of C99 releases the amyloid beta proteins, amyloid-beta 40 (Abeta40) and amyloid-beta 42 (Abeta42), major components of amyloid plaques, and the cytotoxic C-terminal fragments, gamma-CTF(50), gamma-CTF(57) and gamma-CTF(59).
Host / Isotype:	Mouse / IgG1
Recommended Isotype Controls:	SM10B (for use in human samples)
Clone:	19H11
Immunogen:	N-terminal peptide conjugated to KLH. Epitope: N-Terminus of Amyloid beta-A4: NH2 - D A E F R . .
Format:	State: Liquid purified IgG fraction. Purification: Subsequent Thiophilic Adsorption and Size Exclusion Chromatography Buffer System: 2 x PBS containing 0.09% Sodium Azide, PEG and Sucrose. Label: Biotin
Applications:	ELISA: Use at 0.05 µg/ml. Immunoblotting: 1 µg/ml for HRPO/ECL detection <i>Recommended blocking buffer:</i> Casein/Tween 20 based blocking and blot incubation buffer. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognizes the free N-Terminus of the bA4 polypeptide with high preference and shows only minor crossreactions with APP. Species: Human. Other species not tested.
Storage:	Store the antibody (aliquote and freeze in liquid nitrogen) at -80°C. Thaw aliquots at 37°C. Thawed aliquots may be stored at 4°C up to 3 months. Avoid repeated freezing and thawing. Shelf life: one year from despatch.