

**AM08099PU-N****Monoclonal Antibody to Rabbit Ig (IgG + IgM) - Purified**

<b>Quantity:</b>	0.5 mg
<b>Concentration:</b>	0.5 mg/ml
<b>Host / Isotype:</b>	Rat / IgG2a
<b>Recommended Isotype Controls:</b>	SM15P, SM15PX
<b>Clone:</b>	SB87a
<b>Immunogen:</b>	Rabbit IgG and IgM
<b>Format:</b>	<b>State:</b> Liquid purified Ig fraction. <b>Buffer System:</b> 100 mM Borate Buffered Saline, pH 8.2. No preservatives or amine-containing buffer salts added.
<b>Applications:</b>	<b>ELISA:</b> 1/4,000-1/8,000. <b>Flow Cytometry.</b> Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody reacts with Rabbit IgG and Rabbit IgM as demonstrated by ELISA. It may also react with Ig from other species. <b>Species:</b> Rabbit. Other species not tested.
<b>Storage:</b>	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
<b>Product Citations:</b>	<b>Originator or purchased from resellers:</b> 1. Sang Y, Zhou T, Li H, Ran Y, Jiang D, Zheng F, et al. A new long-acting GLP-1 derivative KTP ameliorates hyperglycemia and dyslipidemia and improves pancreas and fatty liver in db/db mice. Chin Sci Bull. 2013;58:2447-53. 2. Fernando J, Faber TW, Pullen NA, Falanga YT, Kolawole EM, Oskeritzian CA, et al. Genotype-dependent effects of TGF- $\beta$ 1 on mast cell function: targeting the Stat5 pathway. J Immunol. 2013 Nov 1;191(9):4505-13. doi: 10.4049/jimmunol.1202723. Epub 2013 Sep 25. PubMed PMID: 24068671.