

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com **OriGene Technologies GmbH** 

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## AM08120FC-N Monoclonal Antibody to CD4 - FITC


Alternate names:	T-cell surface antigen T4/Leu-3, T-cell surface glycoprotein CD4
Quantity:	0.5 mg
Concentration:	0.5 mg/ml
Background:	CD4 is a single chain transmembraneous glycoprotein (59 kDa) which belongs to the immunoglobulin superfamily. CD4 is present on a subset of T lymphocytes ("helper/inducer" T cells) and is also expressed at a lower level on monocytes, tissue macrophages and granulocytes. The antigen is involved in binding to MHC class II molecules. The intracellular domain of the antigen is associated with p56lck protein tyrosine kinase.
Uniprot ID:	<u>Q6Q4G0</u>
NCBI:	<u>9031</u>
Host / Isotype:	Mouse / IgM
Recommended Isotype Controls:	SM13F
Clone:	EP96
Format:	<b>State:</b> Liquid purified Ig fraction. <b>Buffer System:</b> PBS containing 0.09% Sodium Azide as preservative. <b>Label:</b> FITC – Fluorescein Isothiocyanate Isomer 1
Applications:	<b>Flow Cytometry:</b> $\langle / = 1 \ \mu g / 10e6$ cells. (Ref.1) Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognizes Chjicken CD4 (Mr. 64 kDa). EP96 inhibits both PHA- and Con A-induced proliferative responses of splenocytes, and PWM-induced IL-2 production. (Ref.1) <b>Species:</b> Chicken. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for longer. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	1. Ratcliffe, M. 1996. Personal communication. 2. Cooper, M.D., R.P. Bucy, and CL. Chen. 1990. In: The Avian Model in Developmental Biology: From Organism to Genes. Editions du CNRS, pp. 239-249.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

## AM08120FC-N: Monoclonal Antibody to CD4 - FITC

## **Pictures:**

Immunofluorescent Staining: Chicken peripheral blood mononuclear cells were double stained with anti-Chicken CD4-FITC and anti-Chicken TCR alpha/beta-PE (AM08114FC-N). Small lymphocytes were then gated and analyzed on a FACScan(TM) flow cytometer (BDIS, San Jose, CA). Amount Used: 1 µg/10e6 cells.



For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.