

AM08124PU-N**Monoclonal Antibody to MHC Class II (I-A) - Purified**

Quantity:	0.5 mg
Concentration:	0.5 mg/ml
Background:	The Chicken Ia antigen is expressed on B cells, a subpopulation of the monocyte/macrophage lineage of cells, and mitogen-activated T cells. (Ref.1)
Host / Isotype:	Mouse / IgM
Recommended Isotype Controls:	SM13P
Clone:	Cla-I
Format:	State: Liquid purified Ig fraction. Buffer System: 100 mM Borate Buffered Saline, pH 8.2. No preservatives or amine-containing buffer salts added.
Applications:	Flow Cytometry: Identification and enumeration of Ia+ cells. (Ref.1) Immunoprecipitation. (Ref.1) Inhibition of MLR. (Ref.1) Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognizes a 65 kDa glycoprotein consisting of a 32 kDa alpha-chain non-covalently bound to a 27 kDa beta-chain. The molecule recognized by this antibody is similar in structure to the murine Ia antigens and human DR antigens. Species: Chicken. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	1. Ewert, D.L., M.S. Munchus, C.-L. Chen, and M.D. Cooper. 1984. J. Immunol. 132:2524.