

AM08129PU-N**Monoclonal Antibody to Chicken Lambda light chain - Purified**

Alternate names:	Ig lambda chain C regions
Quantity:	0.5 mg
Concentration:	0.5 mg/ml
Background:	All five immunoglobulin classes share the same basic four polypeptide chain structure of two heavy-chains and two light chains. There are five heavy chain types, and two light-chain types (Kappa and Lambda) both having a molecular weight of 22.5kDa. Any heavy-chain type can associate with either light-chain type, but on any immunoglobulin molecule both light-chains are of the same type. Kappa and Lambda consist of a variable region and a constant region and can easily be differentiated by the antigenic properties of the constant region. The ratio of Kappa to Lambda is 70:30, the vast majority of which is bound to heavy-chain in immunoglobulin. In normal individuals low levels of free light-chain are present in serum (kappa, 1.6-15.2 mg/L; Lambda, 0.4-4.2mg/L), with the occurrence of multiple myeloma or other B-cell malignancies these levels can be greatly elevated and can be found at high levels in the urine (Bence-Jones proteins).
Host / Isotype:	Mouse / IgG1
Recommended Isotype Controls:	AM03095PU-N
Clone:	L-1
Format:	State: Liquid purified Ig fraction. Buffer System: 100 mM Borate Buffered Saline, pH 8.2. No preservatives or amine-containing buffer salts added.
Applications:	ELISA. Flow Cytometry. Immunofluorescence. Immunohistochemistry on Frozen Sections. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody recognizes Lambda light chains of Chicken Immunoglobulin. (Ref.1) Species: Chicken. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
General Readings:	1. Chen, C.H., J.E. Lehmeyer, and M.D. Cooper. 1982. J. Immunol. 129:2580-2585.