

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AM08141FC-N **Monoclonal Antibody to CD8 - FITC**

Alternate names: CD8 alpha chain, CD8A, MAL, T-cell surface glycoprotein CD8 alpha chain, T-

lymphocyte differentiation antigen T8/Leu-2

0.5 mg **Quantity: Concentration:** 0.5 mg/ml

Background: The CD8 antigen is a cell surface glycoprotein found on most cytotoxic Tlymphocytes

that mediates efficient cell to cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T cell receptor on the T lymphocyte recognize antigen displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. CD8 is expressed on most thymocytes and approximately 1/3 of peripheral blood T cells in humans. CD8 alpha/beta heterodimers are expressed only on TCR alpha/beta T cells, whereas CD8 alpha homodimers can be expressed on alpha/beta and gamma/delta T cells and some NK cells in humans. The pattern of expression in other species has not

been well documented.

Uniprot ID: 00R4I4 NCBI: 9031

Host / Isotype: Mouse / IgG2b

Recommended Isotype

Controls: Clone:

SM12F

3-298

Format:

State: Liquid purified Ig fraction.

Buffer System: PBS containing 0.09% Sodium Azide as preservative.

Label: FITC – Fluorescein Isothiocyanate Isomer 1

Applications: Flow Cytometry: $\langle / = 1 \mu g / 10e6 \text{ cells.}$ (Ref.4)

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

This antibody recognizes the CD8 alpha chain of both the Chicken and Turkey. (Ref.4) **Specificity:**

> **Species:** Chicken and Turkey. Other species not tested.

Store the antibody undiluted at 2-8°C for one month or in (aliquots) at -20°C for Storage:

longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings: 1. Chan, M.L., C.-L. Chen, L. Ager, and M.D. Cooper. 1988. J. Immunol. 140:2133-2138.

2. Cooper, M.D., R.P. Bucy, and C.-L. Chen. 1990. In: The Avian Model in



Developmental Biology: From Organism to Genes. Editions du CNRS, pp. 239-249.

- 3. Tregaskes, C.A., F. Kong, E. Paramithiotis, C.-L. Chen, M.J.H. Ratcliffe, T. F. Davison, and J.R. Young. 1995. J. Immunol. 154:4485-4494.
- 4. Li, Z., K.E. Nestor, Y.M. Saif, Z. Fan, M. Luhtala, and O. Vainio. 1999. Poult Sci. 78:1526-1531.