

OriGene Technologies Inc.

AM08398FC-N
OriGene EU

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 <u>US-info@acris-antibodies.com</u>

Monoclonal Antibody to CD45 - FITC

Alternate names: CD45R, CD45RA, CD45RB, CD45RO, L-CA, Leukocyte common antigen, PTPRC, T200

Catalog No.: AM08398FC-N

Quantity: 50 μg

Background: CD45 is a family of single chain transmembraneous glycoproteins consisting of at least four

isoforms (220, 205, 190, 180 kDa) which share a common large intracellular domain. Their extracellular domains are heavily glycosylated. The different isoforms are produced by alternative messenger RNA splicing of three exons of a single gene on chromosome 1. CD45 is expressed on cells of the human hematopoietic lineage (including hematopoietic stem cells) with the exception of mature red cells. It is not detected on differentiated cells of other tissues. It is likely that CD45 plays an important role in signal transduction, inhibition or upregulation of various immunological functions. Antibodies recognising a common epitope on all of the isoforms are termed CD45 whilst those recognising only individual

isoforms are termed CD45RA or CD45RO etc.

Host / Isotype: Mouse / IgG1

Clone: 1H-1

Format: State: Liquid purified lg fraction.

Purification: Protein A Chromatography.

Buffer System: PBS, pH 8.0 containing 1% BSA as stabilizer and 0.09% Sodium Azide as

preservative.

Label: FITC - Fluorescein

Applications: Flow Cytometry: Use 10 µl of neat-1/10 diluted antobody to label 10e6 cells in 100 µl.

Immunohistochemistry on Paraffin Sections: 10 μg/ml.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: AM08398FC-N recognizes CD45 on all leucocytes.

Immunoprecipitates proteins of M.Wt. 220,000, 210,000, 195,000 and 180,000 from lymph node lymphocytes and proteins of M.Wt 195,000, 180,000 and 170,000 from thymocytes.

Species Reactivity: Tested: Guinea Pig.

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This Product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings: 1. Hart IJ, Schäfer H, Scheper RJ, Stevenson GT. Subpopulations of guinea-pig T

lymphocytes defined by isoforms of the leucocyte common antigen. Immunology. 1992

Nov;77(3):377-84. PubMed PMID: 1362184.

TEV NORD
TOWNORD CERT
OFFI



AM08398FC-N: Monoclonal Antibody to CD45 - FITC

Pictures:

Spleen: Formalin-Fixed Paraffin-Embedded (FFPE)

