

OriGene Technologies Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES

Phone: +1-858-888-7900 Fax: +1-858-888-7904 US-info@acris-antibodies.com

AM10216SU-N OriGene EU

Acris Antibodies GmbH

Schillerstr. 5 32052 Herford GERMANY

Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info@acris-antibodies.com

Monoclonal Antibody to Rotavirus VP6 protein - Ascites

Catalog No.: AM10216SU-N

Quantity: 0.2 ml

Background: Rotaviruses, members of the family Reoviridae, are a major cause of diarrhoea in young

mammals. Rotavirus infections also result in economic losses in agriculture due to diarrhoea in calf, pig, sheep, and poultry rearing. Diarrhoea (or scours) due to the rotavirus Nebraska Calf Diarrhea Virus can affect calves up to 30 days of age or older. Diarrhoea begins 2 to 3 days after exposure. Diagnosis is by history, lesions (ulcers on the tongue, lips, and mouth) and diagnostic laboratory tests. Mortality rates may be as high as 50 percent, depending on the secondary bacteria present. Human rotaviruses, the major aetiological agents of severe infantile diarrhoea worldwide, display surprisingly diverse and complex serotypic specificities. Rotaviruses are 70 nm, non enveloped viruses comprised of a triple layered protein capsid; Outer capsid proteins are VP4 and VP7, Inner capsid -VP6 and Core -VP2. The immunity acquired from exposure to rotavirus appears to be type specific following initial infection; therefore, multiple serotypes of rotavirus mean multiple opportunities for infection. The combination of animal reservoirs for the virus and rotavirus gene reassortment provides the potential for dramatic genetic shifts (similar to

influenza virus) which could give rise to altered host ranges and viral virulence.

Host / Isotype: Mouse / IgG2b

Clone: 2B4

Immunogen: Recombinant inner capsid protein (VP6)

Format: State: Lyophilized powder

Preservatives: None

Reconstitution: Restore in distilled water.

Applications: Western Blot: 1/200-1/1000.

Immunocytochemistry/Immunofluorescence: 1/100-1/500.

Immunohistochemistry: 1/100-1/500.

Other applications not tested. Optimal dilutions are dependent on conditions and should

be determined by the user.

Specificity: Reacts with Human 41 kDa inner capsid protein (VP6) of Rotavirus (RV).

Storage: Prior to reconstitution store the antibody at -20°C.

Store reconstituted antibody at 2-8°C for one month or (in aliquots) at -20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

