

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AM26153PU-N Monoclonal Antibody to CHIPS (N-term) - Purified

Quantity: 0.1 mg
Concentration: 0.1 mg/ml

Background: The bacterial pathogen Staphylococcus aureus is insensitive to antimicrobial host

defense peptides such as defensins, protegrins, platelet microbicidal proteins and

bacteriocins. Staphylococci have developed various resistance mechanisms

including those specific for bacteriocins and several host defense peptides. A protein belonging to the resistance mechanism of Staphylococcus aureus is known as CHIPS

(Chemotaxis Inhibiting Protein of Staphylococcus aureus). CHIPS is a protein produced by Staphylococcus aureus that inhibits chemotaxis of neutrophils by blocking the Formyl Peptide Receptor (FPR) and C5a Receptor on neutrophils. CHIPS and antibodies against CHIPS can be useful for various experimental infection models of Staphylococcus aureus. Furthermore these reagents can be of help in studies on

the role of FPR and C5a in inflammatory processes.

Uniprot ID: Q6GFB3

NCBI: <u>YP_041409.1</u>

GeneID: 2861356

Host / Isotype: Mouse / IgG1

Recommended Isotype AM03095PU-N

Controls:

Clone: INC1

Format: State: Liquid 0.2 µm filtered Ig fraction

Purification: Protein G purified

Buffer System: PBS

Preservatives: 0.01% sodium azide

Stabilizers: 0.1% BSA

Applications: Immuno assays.

Flow cytometry (1:10 as starting dilution). Western blot (1:100 as starting dilution).

Immunohistology on frozen sections (1:100 as starting dilution). Immunohistology on paraffin sections (1:100 as starting dilution).

Inhibition of biological activity.

Before use in biological assays, the product must be filter sterilized and depending on the concentration to be used dialyzed against culture medium to remove the

sodium azide added.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: This antibody detects reacts the N-terminus of CHIPS.

Storage: Product should be stored at 2-8°C. Under recommended storage conditions, product

is stable for one year.