

AM32826PU-S Monoclonal Antibody to Plasma Cell - Purified

Alternate names:	Plasma B cell, Plasmocytes
Quantity:	0.1 mg
Concentration:	0.2 mg/ml
Background:	Plasma cells (also called plasma B cells or plasmocytes), which are large lymphocytes derived from an antigen-specific B cell, secrete antibodies and are responsible for humoral immunity. Plasma cells differentiate from B cells upon stimulation by CD4+ lymphocytes. The B cell acts as an antigen-presenting cell (APC), consuming an offending pathogen, which is taken up by the B cell by phagocytosis and broken down within proteosomes. Plasma cells contain basophilic cytoplasm; their nucleus contains heterochromatin organized in a characteristic cartwheel arrangement.
Host / Isotype:	Mouse / IgG2a
Recommended Isotype Controls:	AM03096PU-N
Clone:	LIV3G11
Immunogen:	Pancreatic cancer related serum Mucin.
Format:	State: Liquid purified IgG fraction from Bioreactor Concentrate Purification: Protein A/G Chromatography Buffer System: 10mM PBS Preservatives: 0.05% Sodium Azide Stabilizers: 0.05% BSA
Applications:	Immunohistochemistry on Paraffin Sections: Use Plasma Cell antibody at 0.5-1 µg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. Positive Control: Tonsil or Lymph node. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	LIV3G11 antibody recognizes an intracytoplasmic antigen which shows a very high degree of specificity for plasma cells. This antigen is present in normal as well as neoplastic plasma cells. This monoclonal antibody superbly recognizes normal and neoplastic plasma cells in routine formalin-fixed, paraffin-embedded tissue sections. Plasma Cell Marker antibody is of potential value in identifying myeloma or plasmacytoma in bone marrow or other tissues. It also helps differentiate lymphoplasmacytoid lymphoma from lymphocytic and follicular lymphoma. Note that this MAb is not suitable for staining frozen tissues. Cellular Localization: Cytoplasmic. Negative Species: Rat.
Species Reactivity:	Tested: Human.

Add. Information: Clone LIV3G11 is the same as 7B18

Storage: Store undiluted at 2-8°C.
Shelf life: one year from despatch.

General Readings:

1. Turley H, Jones M, Erber W, Mayne K, de Waele M, Gatter K. VS38: a new monoclonal antibody for detecting plasma cell differentiation in routine sections. *J Clin Pathol.* 1994 May;47(5):418-22. PubMed PMID: 7517959.
2. Ching CK and Rhodes JM. 1998. *Gastroenterology*; 95: 137-42.
3. Ching CK, Rhodes JM. Purification and characterization of a peanut-agglutinin-binding pancreatic-cancer-related serum mucus glycoprotein. *Int J Cancer.* 1990 Jun 15;45(6):1022-7. PubMed PMID: 2351483.

Pictures: Formalin-paraffin human tonsil stained with Plasma Cell Marker MAb (Clone LIV3G11).

