

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AM33178SU-N Polyclonal Antibody to CD254 / RANKL (287-304) - Serum

Alternate names: ODF, OPGL, Osteoclast differentiation factor, Osteoprotegerin ligand, RANK Ligand,

RANKL, Receptor activator of nuclear factor kappa B ligand, TNF-related activation-induced cytokine, TNFSF11, TRANCE, Tumor necrosis factor ligand superfamily member

11

Quantity: 0.1 ml

Background: RANKL is a member of the tumor necrosis factor (TNF) cytokine family which is a ligand

for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. There are three isoforms of RANKL. Human RANKL is a soluble 20 kDa polypeptide, comprising the TNF homologous region of RANKL (176 amino acid residues). This protein was shown to be a dentritic cell survival factor and is involved in the regulation of T cell dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor associated factor (TRAF) 6, which indicated that this protein may have a role in the regulation of cell apoptosis. RANKL deficient mice show severe osteoporesis and

complete absence of osteoclasts as a result of lack of osteogenesis.

Uniprot ID: <u>014788</u>

NCBI: NP 003692

GeneID: 8600
Host: Rabbit

Immunogen: Human sRANKL synthetic Peptide (EEISIEVSNPSLLDPDQD)

Genename: TNFSF11

Format: State: Lyophilized Serum.

Reconstitution: Restore in aqua bidest to initial volume.

Applications: ELISA.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: Recognizes Human sRANKL (soluble form). Other species not tested.

Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month

or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing. Shelf life: one year from despatch.