

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## **OriGene Technologies GmbH**

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## AM33178SU-S Polyclonal Antibody to CD254 / RANKL (287-304) - Serum

Alternate names:	ODF, OPGL, Osteoclast differentiation factor, Osteoprotegerin ligand, RANK Ligand, RANKL, Receptor activator of nuclear factor kappa B ligand, TNF-related activation- induced cytokine, TNFSF11, TRANCE, Tumor necrosis factor ligand superfamily member 11
Quantity:	20 µl
Background:	RANKL is a member of the tumor necrosis factor (TNF) cytokine family which is a ligand for osteoprotegerin and functions as a key factor for osteoclast differentiation and activation. There are three isoforms of RANKL. Human RANKL is a soluble 20 kDa polypeptide, comprising the TNF homologous region of RANKL (176 amino acid residues). This protein was shown to be a dentritic cell survival factor and is involved in the regulation of T cell dependent immune response. T cell activation was reported to induce expression of this gene and lead to an increase of osteoclastogenesis and bone loss. This protein was shown to activate antiapoptotic kinase AKT/PKB through a signaling complex involving SRC kinase and tumor necrosis factor receptor associated factor (TRAF) 6, which indicated that this protein may have a role in the regulation of cell apoptosis. RANKL deficient mice show severe osteoporesis and complete absence of osteoclasts as a result of lack of osteogenesis.
Uniprot ID:	<u>014788</u>
NCBI:	<u>NP_003692</u>
GenelD:	<u>8600</u>
Host:	Rabbit
Immunogen:	Human sRANKL synthetic Peptide (EEISIEVSNPSLLDPDQD) Genename: TNFSF11
Format:	<b>State:</b> Lyophilized Serum. <b>Reconstitution:</b> Restore in aqua bidest to initial volume.
Applications:	<b>ELISA.</b> Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	Recognizes Human sRANKL (soluble form). Other species not tested.
Storage:	Store lyophilized at 2-8°C for 6 months or at -20°C long term. After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C long term. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

MS/20140325 1/1