

AM33324PU-N**Monoclonal Antibody to Human IgM (heavy chain) - Purified**

Alternate names:	Human Immunoglobulin M
Quantity:	0.2 mg
Concentration:	0.2 mg/ml
Host / Isotype:	Mouse / IgG2b
Recommended Isotype Controls:	SM12P, AM03110PU-N
Clone:	IM373
Immunogen:	Recombinant heavy chain of Human IgM. Genename: IGHM
Format:	State: Liquid purified IgG fraction from Bioreactor Concentrate Purification: Protein A/G Chromatography Buffer System: 10mM PBS Preservatives: 0.05% Sodium Azide Stabilizers: 0.05% BSA
Applications:	ELISA: Use Antibody without BSA for Coating. Flow Cytometry: 0.5-1 µg/10 ⁶ cells. Immunofluorescence: 0.5-1 µg/ml. Western Blot: 0.5-1 µg/ml. Immunoprecipitation: 0.5-1 µg/500 µg protein lysate. Immunohistochemistry on Frozen and Formalin-Fixed Sections: 0.5-1 µg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. Recommended Positive Control: 293T, Raji or hPBL cells, Tonsil or Spleen. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Molecular Weight:	50-75kDa
Specificity:	Recognizes a protein of 75kDa, identified as mu heavy chain of Human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This Monoclonal Antibody is useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin's lymphomas. The most common feature of these malignancies is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant. Cellular Localization: Cytoplasm, Cell Surface and Secreted. Species: Human. Other species not tested.

Storage: Store undiluted at 2-8°C.
Shelf life: one year from despatch.

General Readings: 1. Baryshnikov Alu. [Monoclonal antibodies of the ICO series against differentiation antigens of human lymphocytes]. Gematol Transfuziol. 1990 Aug;35(8):4-7. PubMed PMID: 2253863.

Pictures: Formalin-paraffin human tonsil stained with IgM MAb (Clone IM373)

