

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AM33338PU-S Monoclonal Antibody to Nuclear Antigen (Membrane) - Purified

Alternate names: Nuclear Membrane Marker

Quantity: 0.1 mg
Concentration: 0.2 mg/ml

Background: The nuclear envelope is riddled with nuclear pores that allow specific materials to

pass in and out of the nucleus. Attached to the nuclear envelope is the endoplasmic

reticulum.

Host / Isotype: Mouse / IgG1

Recommended Isotype

Controls:

SM10P (for use in human samples), AM03095PU-N

Clone: NM97

Immunogen: Nuclei of myeloid leukemia biopsy cells.

Format: State: Liquid purified IgG fraction from Bioreactor Concentrate

Purification: Protein A/G Chromatography

Buffer System: 10mM PBS

Preservatives: 0.05% Sodium Azide

Stabilizers: 0.05% BSA

Applications: Immunofluorescence: 0.5-1 μg/ml.

Immunocytochemistry (Acetone-fixed cells): 0.5-1.0 μg/ml for 30 minutes at RT. Immunohistochemistry on Frozen and Formalin-Paraffin Sections: 0.5-1 μg/ml for 30

minutes at RT.

Recommended Positive Control: Tonsil.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: This Monoclonal Antibody is part of a new panel of rea gents, which recognizes

subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This Monoclonal NM97 Antibody recognizes an antigen associated with the nuclear membrane express ed in human cells. It can be used to stain the nuclear membrane in cell or tissue preparations and can be used as a marker of the nuclear membrane in subcellular fractions. It produces a ring pattern around the nucleus of cells of normal and malignant cells and may be used to stain the nuclear membrane of cells in fixed or frozen tissue sections. It can be used with paraformaldehyde fixed frozen tissue or

cell preparations and formalin fixed, paraffin -embedded tissue sections.

Cellular Localization: Nuclear membrane.

Species Reactivity: Tested: Human.

Storage: Store undiluted at 2-8°C.

DO NOT FREEZE!

Shelf life: one year from despatch.