

**AM33339PU-S****Monoclonal Antibody to Nucleoli Marker - Purified**

<b>Quantity:</b>	0.1 mg
<b>Concentration:</b>	0.2 mg/ml
<b>Host / Isotype:</b>	Mouse / IgG1
<b>Recommended Isotype Controls:</b>	SM10P (for use in human samples), AM03095PU-N
<b>Clone:</b>	NM95
<b>Immunogen:</b>	Nuclei of myeloid leukemia biopsy cells.
<b>Format:</b>	<b>State:</b> Liquid purified IgG fraction from Bioreactor Concentrate <b>Purification:</b> Protein A/G Chromatography <b>Buffer System:</b> 10mM PBS <b>Preservatives:</b> 0.05% Sodium Azide <b>Stabilizers:</b> 0.05% BSA
<b>Applications:</b>	<b>Immunofluorescence:</b> 0.5-1 µg/ml. <b>Immunocytochemistry (Acetone-fixed cells):</b> 0.5-1.0 µg/ml for 30 minutes at RT. <b>Immunohistochemistry on Frozen and Paraffin Sections:</b> 0.5-1 µg/ml for 30 minutes at RT. <b><i>Recommended Positive Control:</i></b> Tonsil. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. Monoclonal NM95 Antibody recognizes an antigen associated with the nucleoli in human cells. It can be used to stain the nucleoli in cell or tissue preparations and can be used as a marker of the nucleoli in subcellular fractions. It produces a speckled pattern in the nuclei of cells of normal and malignant cells and may be used to stain the nucleoli of cells in fixed or frozen tissue sections. It can be used with paraformaldehyde fixed frozen tissue or cell preparations and formalin fixed, paraffin-embedded tissue sections. <b><i>Cellular Localization:</i></b> Nucleoli.
<b>Species Reactivity:</b>	<b>Tested:</b> Human.
<b>Storage:</b>	Store undiluted at 2-8°C. <b>DO NOT FREEZE!</b> Shelf life: one year from despatch.