



9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AM50118PU-T Monoclonal Antibody to Human IgG (heavy chain) - Purified


Alternate names:	Human Immunoglobulin G
Quantity:	20 µg
Concentration:	0.2 mg/ml
Host / Isotype:	Mouse / IgG2a
Recommended Isotype Controls:	AM03096PU-N
Clone:	SPM556
Immunogen:	Purified Human Ig Gamma Chain. Genename: IGHG
Format:	State: Liquid purified IgG fraction from Bioreactor Concentrate Purification: Protein A/G Chromatography Buffer System: 10mM PBS Preservatives: 0.05% Sodium Azide Stabilizers: 0.05% BSA
Applications:	 ELISA: Use Antibody without BSA for Coating. Flow Cytometry: 0.5-1 μg/10⁶ cells. Immunofluorescence: 0.5-1 μg/ml. Western Blot: 0.5-1 μg/ml. Immunoprecipitation: 0.5-1 μg/500 μg protein lysate. Immunohistochemistry on Frozen and Formalin-Fixed Paraffin Sections: 0.5-1 μg/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes. <i>Positive Control</i>: 293T, Raji or hPBL cells. Tonsil or Spleen. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Molecular Weight:	75kDa
Specificity:	Recognizes a protein of 75kDa, identified as gamma heavy chain of Human immunoglobulins. It does not cross-react with alpha (IgA), mu (IgM), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is useful in the identification of leukemias, plasmacytomas, and certain non- Hodgkin's lymphomas. The most common feature of these malignancies is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates that the infiltrate is clonal and therefore malignant. <i>Cellular Localization</i> : Cytoplasm, Cell Surface and Secreted. Species: Human. Other species not tested.
Storage:	Store undiluted at 2-8°C. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.



Pictures:

Formalin-Fixed, Paraffin-Embedded Human Tonsil stained with Human IgG Antibody Cat.-No AM50118PU-N (Clone SPM556).



For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.