

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11

info-de@origene.com

AM60041FC-N Monoclonal Antibody to SIGLEC5 / SIGLEC14 - FITC

Alternate names: CD170, O15389, Q08ET2, Sialic acid-binding Ig-like lectin 14, Sialic acid-binding Ig-

like lectin 5

Quantity: 0.1 mg
Concentration: 0.1 mg/ml

Background: Siglec-5 shares an almost identical sequence with Siglec-14 within the first two Ig-like

domains, indicating partial gene conversion between these two Siglecs, also evident

in other primate species.

Siglec-5, also known as CD170, is a sialic-acid-binding Ig-like lectin, and member of the Ig superfamily, expressed by dendritic cells (DCs), activated macrophages, neutrophils, and cells of the monocyte/myeloid lineage. Highly related to the

myelomonocytic-derived adhesion molecule CD33 (Siglec-3), Siglec-5 mediates sialic-acid dependent binding to cells, and is as well acting as an inhibitory receptor in the down-regulation of cell activation. Structurally, Siglec-5 contains an immunoreceptor tyrosine-based inhibitor motif (ITIM), which plays a part in the modulation of cellular responses, and when phosphorylated, can bind to the SH2 domain of several

SH2-containing phosphatases.

Siglec-14 is a putative sialic-acid binding adhesion molecule and predominantly expressed in hematopoietic tissues, which have been shown to associate with the

activating adapter protein DAP12.

Host / Isotype: Mouse / IgG1

Recommended Isotype

Controls:

SM10F (for use in human samples), AM03095PU-N

Clone: 1A5

Immunogen: SIGLEC5-Fc protein, consisting of the full-length extracellular region of human

SIGLEC5, fused with the Fc region of human IgG1

Format: State: Liquid purified IgG

Purification: Affinity chromatography on Protein G

Buffer System: PBS containing 0.09% Sodium azide and 1% BSA

Label: FITC – Fluorescein Isothiocyanate Isomer 1

Applications: Flow cytometry: Neat-1/10, use 10µl of this working dilution to label 10e6 cells in

100µl.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: This antibody recognises human SIGLEC5 / CD170. Clone 1A5 antibody is one of

several SIGLEC5 antibodies which also recognises human SIGLEC14 (Angata, T. et al.,

2006).

Species Reactivity: Tested: Human

Expected from sequence similarity: Chimpanzee (Nguyen, D.H. et al., 2006))

Storage:

Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer.

This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.

General Readings:

1. Cornish AL, Freeman S, Forbes G, Ni J, Zhang M, Cepeda M, et al. Characterization of siglec-5, a novel glycoprotein expressed on myeloid cells related to CD33. Blood. 1998 Sep 15;92(6):2123-32. PubMed PMID: 9731071.

2. Avril T, Freeman SD, Attrill H, Clarke RG, Crocker PR. Siglec-5 (CD170) can mediate inhibitory signaling in the absence of immunoreceptor tyrosine-based inhibitory motif phosphorylation. J Biol Chem. 2005 May 20;280(20):19843-51. Epub 2005 Mar 15. PubMed PMID: 15769739.

3. Nguyen DH, Hurtado-Ziola N, Gagneux P, Varki A. Loss of Siglec expression on T lymphocytes during human evolution. Proc Natl Acad Sci U S A. 2006 May 16;103(20):7765-70. Epub 2006 May 8. PubMed PMID: 16682635.

4. Jaroenpool J, Rogers KA, Pattanapanyasat K, Villinger F, Onlamoon N, Crocker PR, et al. Differences in the constitutive and SIV infection induced expression of Siglecs by hematopoietic cells from non-human primates. Cell Immunol. 2007 Nov-

Dec;250(1-2):91-104. doi: 10.1016/j.cellimm.2008.01.009. Epub 2008 Mar 10. PubMed PMID: 18331725.

5. Angata T, Hayakawa T, Yamanaka M, Varki A, Nakamura M. Discovery of Siglec-14, a novel sialic acid receptor undergoing concerted evolution with Siglec-5 in primates. FASEB J. 2006 Oct;20(12):1964-73. PubMed PMID: 17012248.

Pictures:

Human peripheral blood monocytes stained with SIGLEC5 / SIGLEC14 antibody Cat.-No. AM60041FC-N.

