

AP00003CP-N**Caspase-3 Control Peptide****Alternate names:**

Apopain, CASP-3, CASP3, CPP-32, CPP32, Cysteine protease CPP32, SCA-1, SCA1, SREBP cleavage activity 1, Yama protein

Quantity:

50 µg

Background:

The caspase family of cysteine proteases play a key role in apoptosis. Caspase 3 (also known as CPP32, YAMA and apopain) is the most extensively studied apoptotic protein among caspase family members. Caspase 3 is synthesized as inactive pro enzyme that is processed in cells undergoing apoptosis by self proteolysis and/or cleavage by other upstream proteases (e.g. Caspases 8, 9 and 10).

The processed form of Caspase 3 consists of large (17kD) and small (12kD) subunits which associate to form an active enzyme. Caspase 3 is cleaved at Asp28 - Ser29 and Asp175 - Ser176. The active Caspase 3 proteolytically cleaves and activates other caspases (e.g. Caspases 6, 7 and 9), as well as relevant targets in the cells (e.g. PARP and DFF). Alternative splicing of this gene results in two transcript variants which encode the same protein.

In immunohistochemical studies Caspase 3 expression has been shown to be widespread but not present in all cell types (e.g. commonly reported in epithelial cells of skin, renal proximal tubules and collecting ducts). Differences in the level of Caspase 3 have been reported in cells of short lived nature (eg germinal centre B cells) and those that are long lived (eg mantle zone B cells).

Caspase 3 is the predominant caspase involved in the cleavage of amyloid beta 4A precursor protein, which is associated with neuronal death in Alzheimer's disease.

Uniprot ID:

[P42574](#)

NCBI:

[NP_004337.2](#)

GeneID:

[836](#)

Format:

State: Liquid peptide

Buffer System: PBS, pH 7.2, containing 0.1% BSA and 0.02% sodium azide

Applications:

Western blot: It usually blocks the antibody activity completely in Western blot analysis by incubating the peptide with equal volume of antibody for 30 minutes at 37 °C.

Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.

Specificity:

Control peptide for AP00003PU-N only.

Storage:

Store the antibody undiluted at -20°C or for long term storage (in aliquots) at -70°C.

Avoid repeated freezing and thawing.

Shelf life: one year from despatch.