

AP00107PU-N

Polyclonal Antibody to TANK / ITRAF - Purified

Alternate names:	I-TRAF, TRAF family member-associated NF-kappa-B activator, TRAF-interacting protein, TRAF2
Quantity:	0.1 mg
Concentration:	0.2 mg/ml
Background:	Members of TRAF family associate with activated TNF-R and CD40 and have been implicated to induce nuclear translocation of NFkB. TANK can bind to all three TRAFs and synergizing with TRAF2 to activate the NFkB signal cascade.
Uniprot ID:	Q92844
NCBI:	NP_001186064.1
GeneID:	10010
Host:	Rabbit
Immunogen:	Synthetic peptide surrounding amino acid 413 of human TANK
Format:	State: Liquid purified Ig Purification: Affinity purified Buffer System: PBS pH 7.2, containing 30% glycerol, 0.5% BSA, 0.01% thimerosal.
Applications:	Western Blot: 0.5-4 µg/ml. Immunoprecipitation: 10-20 µg/ml. Immunohistochemistry: 10-20 µg/ml. Mouse small intestine tissue lysate can be used as a positive control. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody detects TANK. Species: Human, Mouse, Rat. Other species not tested.
Storage:	Store the antibody undiluted at -20°C or for long term storage (in aliquots) at -70°C. Avoid repeated freezing and thawing. Shelf life: one year from despatch.
Pictures:	Western blot analysis of TANK with mouse small intestine tissue lysate.

