

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## **OriGene Technologies GmbH**

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## AP01435PU-N Polyclonal Antibody to Vasopressin V2 receptor (AVPR2) - Aff -Purified

Alternate names:	ADHR, AVPR V2, Antidiuretic hormone receptor, DIR, DIR3, Renal-type arginine vasopressin receptor, V2R
Quantity:	0.1 mg
Concentration:	1.0 mg/ml
Background:	Vasopressin (AVP), the antidiuretic hormone, is a cyclic nonpeptide that is involved in the regulation of body fluid osmolality. AVP mediates its effects through a family of G- protein coupled receptors, the vasopressin receptors type V1a, V2 and V3 (also designated V1b). The AVP receptor V1a is responsible for several functions, including blood vessel constriction, liver glycogenolysis and platelet adhesion. It is detected as a full length protein and a shorter protein, which results from proteolytic cleavage of its amino terminus. The V1a receptor is coupled to Gq/11 protein, which increases the intracellular calcium concentration. The human AVP receptor V2 gene maps to chromosome Xq28 and is expressed in lung and kidney. Mutations in the V2 receptor result in nephrogenic diabetes insipidus (NDI), a rare X-linked disorder characterized by the inability of the kidney to concentrate urine in response to AVP. The AVP Receptor V2 activates the Gs protein and the cyclic AMP second messenger system. The AVP Receptor V3 is preferentially expressed in the pituitary and stimulates the release of adrenocorticotropic hormone (ACTH) in response to AVP by mobilizing intracellular calcium stores. AVP receptor antagonists may have potential therapeutic effects in hypertension, congestive heart failure, nephrotic syndrome and ACTH- secreting tumors.
Uniprot ID:	<u>P30518</u>
NCBI:	<u>NP_000045.1</u>
GenelD:	<u>554</u>
Host:	Rabbit
Format:	<ul> <li>State: Liquid purified Ig fraction (&gt; 95% by SDS-PAGE).</li> <li>Purification: Affinity Chromatography using epitope-specific immunogen.</li> <li>Buffer System: Phosphate Buffered Saline (PBS), pH~7.2 with 0.05% Sodium Azide as preservative.</li> </ul>
Applications:	Western Blot: 1/500-1/1000. Immunofluorescence: 1/50-1/200. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Molecular Weight:	~38 kDa

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

	AP01435PU-N: Polyclonal Antibody to Vasopressin V2 receptor (AVPR2) - Aff - Purified		
Specificity:	This antibody detects endogenous levels of AVP Receptor V2 protein. (region surrounding Leu111) <b>Species:</b> Human. Other species not tested.		
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.		
Pictures:	Western blot (WB) analysis of AVP Receptor V2 antibody (CatNo.: AP01435PU-N) in extracts from RAW264.7 cells.		-117 85
		AVP	-49
	Rece	Receptor V2	-34
			-25

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.