

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com **OriGene Technologies GmbH**

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com


AP06242PU-N Polyclonal Antibody to NFKB2 / NF-kappa-B p100/p52 - Aff -Purified

Alternate names:	DNA-binding factor KBF2, H2TF1, KBF2, LYT10, Lymphocyte translocation chromosome 10, NF kappa B, NF-kB p100, NF-kB p52, Nuclear factor NF-kappa-B p100 subunit, Oncogene Lyt-10	
Quantity:	0.1 mg	
Concentration:	1.0 mg/ml	
Background:	The NFkappaB transcription factor was originally identified as a protein complex consisting of a DNA binding subunit and an associated protein. The subunit is functionally related to c-Rel p75 and Rel B p68. The p50 subunit was initially believed to be a functionally unique protein derived from the amino-terminus of a precursor designated p105. A cDNA has been isolated that encodes an alternative DNA binding subunit of NFkappaB. It is synthesized as a protein that is expressed in a variety of cell types and, like p105, undergoes cleavage to generate its NFkappaB subunit, in this case a protein designated p52 (previously referred to as p49). In contrast to p50 derived from p105, p52 acts in synergy with p65 to stimulate the HIV enhancer in transiently transfected Jurkat cells.	
Host:	Rabbit	
Immunogen:	Synthetic peptide, corresponding to amino acids 840-890 of Human NFkB-p100.	
Format:	 State: Liquid purified Ig fraction Purification: Affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE) Buffer System: Phosphate buffered saline (PBS), pH 7.2. Preservatives: 0.05% sodium azide 	
Applications:	Western blot: 1/500-1/1000. Immunohistochemistry on paraffin sections: 1/50-1/200. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.	
Molecular Weight:	~ 100, 120 kDa	
Specificity:	This antibody detects endogenous levels of NFkappaB-p100/p52 protein. (region surrounding Ser869)	
Species Reactivity:	Tested: Human. Expected from sequence similarity: Mouse and Rat.	
Storage:	Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.	

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

	AP06242PU-N: Polyclonal Antibody to NFKB2 / NF-kappa-B p100/p52 - Aff - ORIGENE Purified		
Pictures:	Immunohistochemistry (IHC) analyzes of NFκB-p100 antibody (CatNo.: AP06242PU-N) in paraffin-embedded human breast carcinoma tissue.		

Western blot (WB) analysis of NFkB-p100 antibody (Cat.-No.: AP06242PU-N) in extracts from HeLa cells.



For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.