

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## OriGene Technologies GmbH

Schillerstr. 5 32052 Herford **GERMANY** Phone: +49-5221-34606-0 Fax: +49-5221-34606-11

info-de@origene.com

## Polyclonal Antibody to CHK2 - Aff - Purified AP06380PU-N

CHEK-2, CHEK2, CHK-2, CHK2 checkpoint homolog, Cds1, RAD53, Serine/threonine-Alternate names:

protein kinase Chk2

**Quantity:** 0.1 mg **Concentration:** 1.0 mg/ml

Background: Cell cycle events are regulated by the sequential activation and deactivation of cyclin

dependent kinases (Cdks) and by proteolysis of cyclins. Chk1 and Chk2 are involved in these processes as regulators of Cdks. Chk1 and Chk2 both function as essential components in the G2 DNA damage checkpoint by phosphorylating Cdc25C in response to DNA damage. Phosphorylation inhibits Cdc25C activity, thereby blocking mitosis. Cdc25A, Cdc25B and Cdc25C protein tyrosine phosphatases function as mitotic activators by dephosphorylating Cdc2 p34 on regulatory tyrosine residues. It has also been shown that Chk1 can phosphorylate Wee1 in vitro, providing evidence

that the hyperphosphorylated form of Wee1, seen in cells delayed by Chk1

overexpression, is due to phosphorylation by Chk1.

**Uniprot ID:** 096017

NCBI: NP 001005735.1

GeneID: 11200 Rabbit Host:

Immunogen: Synthetic peptide, corresponding to amino acids 351-400 of Human Chk2.

Format: **State:** Liquid purified Ig fraction

Purification: Affinity-chromatography using epitope-specific immunogen and the

purity is > 95% (by SDS-PAGE)

Buffer System: Phosphate buffered saline (PBS), pH 7.2.

Preservatives: 0.05% sodium azide

**Applications:** Western blot: 1/500-1/1000.

Immunohistochemistry on paraffin sections: 1/50-1/200.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

**Molecular Weight:** ~ 62 kDa

**Specificity:** This antibody detects endogenous levels of Chk2 protein.

(region surrounding Meth381)

**Species Reactivity:** Tested: Human.

Expected from sequence similarity: Mouse and Rat.


Store undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. **Storage:** 

> Avoid repeated freezing and thawing. Shelf life: one year from despatch.



**Pictures:** 

Immunohistochemistry (IHC) analyzes of Chk2 antibody (Cat.-No.: AP06380PU-N) in paraffin-embedded human breast carcinoma tissue.



**Western blot** (WB) analysis of Chk2 antibody (Cat.-No.: AP06380PU-N) in extracts from Jurkat cells.

