

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Polyclonal Antibody to CDK6 - Aff - Purified AP08101PU-N Cell division protein kinase 6, PLSTIRE, Serine/threonine-protein kinase PLSTIRE Alternate names: **Quantity:** 0.1 mg **Concentration:** 1.0 mg/ml **Background:** Cdk6 is a member of the cyclin-dependent protein kinase (CDK) family. CDK family members are highly similar to the gene products of Saccharomyces cerevisiae cdc28, and Schizosaccharomyces pombe cdc2, and are known to be important regulators of cell cycle progression. This kinase is a catalytic subunit of the protein kinase complex that is important for cell cycle G1 phase progression and G1/S transition. The activity of Cdk6 first appears in mid-G1 phase, and is controlled by regulatory subunits including D-type cyclins and members of INK4 family of CDK inhibitors. This kinase, as well as CDK4, has been shown to phosphorylate, and thus regulate the activity of, tumor suppressor protein Rb. **Uniprot ID:** <u>Q00534</u> NCBI: NP 001138778.1 GenelD: <u>1021</u> Host: Rabbit Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human CDK6 around the phosphorylation site of Tyrosine 13 (Q-Q-Yp-E-C). Format: State: Liquid purified lg fraction. **Purification:** Immunoaffinity Chromatography using epitope-specific immunogen. Buffer System: PBS (without Mg2+ and Ca2+), pH 7.4, 150 mM NaCl, 0.02% Sodium Azide and 50% Glycerol. **Applications:** Immunofluorescence: 1/100~1/200. Immunohistochemistry on Paraffin-Embedded Sections: 1/50~1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user. **Specificity:** This antibody detects endogenous levels of total CDK6 protein. Species: Human and Mouse. Other species not tested. Storage: Store the antibody (in aliquots) at -20°C. Avoid repeated freezing and thawing.

 Shelf life: One year from despatch.

 General Readings:
 1. Flores-Rozas, H. et al. (1994) Proc. Natl. Acad. Sci. USA 91, 8655-8659.

 2. Pestell, R.G. et al. (1999) Endocr. Rev. 20, 501-534.
 3. Lukas, J. et al. (1996) Mol. Cell. Biol. 16, 6917-1625.


For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.

OG/20130513 1/2




Pictures:

Figure 1. Immunohistochemical analysis of paraffin-embedded human colon carcinoma tissue using CDK6 Antibody AP08101PU.



Peptide

Figure 2. Immunofluorescence staining of methanol-fixed HeLa cells using CDK6 antibody AP08101PU (Red).



For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.