

AP08136PU-N**Polyclonal Antibody to Luteinizing hormone beta (LSH-beta) - Purified**

Alternate names:	LH-B, LHB, LSH-beta, Luteinizing hormone subunit beta, Lutropin beta chain, Lutropin subunit beta
Quantity:	1 ml
Concentration:	2.0 mg/ml
Background:	Luteinizing Hormone (LH) is a tropic hormone which modulates the secretory activity of other endocrine glands. This glycoprotein hormone is produced in the anterior hypophysis of the pituitary gland. In the female an acute rise of LH the LH surge – triggers ovulation. In the male, where LH had also been called Interstitial Cell Stimulating Hormone (ICSH), it stimulates Leydig cell production of testosterone. The alpha subunits of LH, FSH, TSH, and hCG are identical, and contain 92 amino acids, the beta subunit vary. LH has a beta subunit of 121 amino acids (LHB) that confers its specific biologic action. and is responsible for interaction with the LH receptor.
Uniprot ID:	P01229
NCBI:	NP_000885.1
GeneID:	3972
Host / Isotype:	Rabbit / IgG
Immunogen:	LH beta.
Format:	State: Liquid purified Ig fraction. Purification: Protein A Chromatography. Buffer System: 10 mM PO ₄ , 150 mM NaCl, pH 7.4 containing 0.05% Sodium Azide as preservative and 1% BSA as stabilizer.
Applications:	Suitable for Immunohistochemistry and Immunocytochemistry (Frozen or Formalin-Fixed Paraffin-Embedded (FFPE) tissue sections and cell smears): 1/100-1/500 using streptavidin~biotin system or polymer system. Positive Control: Human anterior pituitary. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody stains both free and intact LH and is useful for the classification of pituitary adenomas and tumors of the pituitary. Cellular Localization: Cytoplasmic. Species: Human. Other species not tested.
Storage:	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: One year from despatch.