

## Polyclonal antibody to Ferritin F(ab)2 fragment - Ig Fraction

<b>Alternate names:</b>	FTH, FTH1, FTHL6, FTL, Ferritin H, Ferritin L
<b>Catalog No.:</b>	AP09202PU-N
<b>Quantity:</b>	5 mg
<b>Concentration:</b>	0.84 mg/ml (by UV absorbance at 280 nm)
<b>Host / Isotype:</b>	Rabbit / IgG
<b>Immunogen:</b>	Ferritin [Human Spleen]
<b>Format:</b>	<b>State:</b> Liquid F(ab)2 fragment of IgG fraction <b>Buffer System:</b> 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 0.01% (w/v) Sodium Azide
<b>Applications:</b>	Suitable for highly specific immunological methods requiring low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This is a IgG F(ab)2 Fragment of anti-Ferritin.
<b>Storage:</b>	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.