

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

AP09203TR-N Polyclonal Antibody to FITC - Texas Red

Alternate names:	Fluorescein Isothiocyanate
Quantity:	1 mg
Concentration:	1.0 mg/ml (by UV absorbance at 280 nm)
Background:	Fluorescein is a fluorophore commonly used to label proteins - protein-fluorescein conjugates are not usually susceptible to precipitation. In addition to its relatively high absorptivity, excellent fluorescence quantum yield and good water solubility, fluorescein has an excitation maximum of 494 nm that closely matches the 488 nm spectral line of the argon-ion laser, making it an important fluorophore for confocal laser-scanning microscopy and flow cytometry applications. Its fluorescence is pH sensitive and is significantly reduced below pH 7. Fluorescein emits most strongly between 500 and 550 nm, but it has a relatively broad emission spectrum reaching to over 600 nm. Several derivatives of fluorescein are commonly used, including FITC (fluorescein isothiocyanate), carboxylates and succinimidyl esters.
Host / Isotype:	Goat / IgG
Immunogen:	Fluorescein conjugated to Goat IgG
Format:	 State: Lyophilized purified Ig Purification: Affinity chromatography Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA) as stabilizer and 0.01% (w/v) Sodium Azide as preservative Label: Texas Red – (TM) Sulfonyl Chloride (Molecular Weight 625 daltons) Absorption / Emission: 596 nm / 620 nm Molar Ratio: 2.9 moles Texas Red(TM) per mole of Goat IgG Reconstitution: Restore with 1,0 ml of deionized water (or equivalent).
Applications:	Flow Cytometry. Immunofluorescence. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody reacts to Fluorescein.
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.