

AP09815PU-N**Polyclonal Antibody to Stanozolol - Ig Fraction**

Alternate names:	Androstanazole, Methylstanazole, Winstrol
Quantity:	0.1 ml
Concentration:	6.6mg/ml (U.V. abs)
Host / Isotype:	Sheep / IgG
Immunogen:	Stanozolol (17)-BSA
Format:	State: Liquid Ig fraction prepared by Caprylic Acid and Ammonium Sulphate precipitation procedures Buffer System: 20mM Phosphate, 150mM Sodium Chloride, pH 7.2 containing 0.09% Sodium Azide as preservative
Applications:	ELISA: 6.6µg/ml. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	This antibody reacts to Stanozolol. Cross-Reactivity; Stanozolol 100% 16 beta-Hydroxystanozolol 40% 4 alpha-Hydroxystanozolol 0.9% 3-Hydroxy-17-Epi-Stanozolol 0.1% 3-Hydroxystanozolol 0.1% 19-Nortestosterone (17alpha) 0.1% 19-Nortestosterone (17beta) 0.1% Trenbolone (17alpha) 0.1% Trenbolone (17beta) 0.1% 19-Nor-4-Androstene - 3,17-Dione 0.1% Boldenone 0.1% Progesterone 0.1% Testosterone (17beta) 0.1% Cortisol 0.1% Estrilol 0.1% Estrone 0.1% Danazol 0.1% Methandriol 0.1% 17alpha-Methyltestosterone 0.1%
Storage:	Store the antibody at -20°C. Avoid repeated freezing and thawing. Shelf life: one year from despatch.