

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606

techsupport@origene.com

OriGene Technologies GmbH

32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Schillerstr. 5

AP20877PU-N Polyclonal Antibody to BRCA1 / RNF53 pSer1524 - Aff - Purified

Alternate names: Breast cancer type 1 susceptibility protein, RING finger protein 53

Quantity: 0.1 mg
Concentration: 1,0 mg/ml

Background: In 1990, a breast cancer susceptibility gene designated BRCA1 was localized to

chromosome 17q. Mutations within this gene are believed to account for

approximately 45% of families with high incidence of breast cancer and at least 80% of families with increased incidence of both early-onset breast cancer and ovarian cancer. A second breast cancer susceptibility gene, BRCA2, located on chromosome 13q12-13, also confers a high incidence of breast cancer but, unlike BRCA1, does not confer a substantially elevated risk of ovarian cancer. The BRCA1 gene is expressed in numerous tissues, including breast and ovary, and encodes a predicted protein of 1863 amino acids. This protein contains a zinc finger domain in its amino terminal region, but is otherwise unrelated to any previously described proteins. Like many other genes involved in familial cancer, BRCA1 appears to encode a tumor suppressor,

a protein that acts as a negative regulator of tumor growth.

 Uniprot ID:
 P38398

 NCBI:
 NP 009225

GeneID: 672
Host: Rabbit

Format: State: Liquid purified lg fraction

Purification: Affinity chromatography (> 95% (by SDS-PAGE)

Buffer System: Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH

7.2.

Applications: Immunohistochemistry: 1/50 - 1/200.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: p-BRCA1 (pSer1524) pAb detects endogenous levels of p-BRCA1 protein

Species: Human.

Other species not tested.

Storage: Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for

onger.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.



Pictures:

Immunohistochemistry (IHC) analysis of p-BRCA1 (pSer1524) pAb in paraffinembedded human breast carcinoma tissue.

