

**AP21261AF-N****Polyclonal Antibody to G6PD - Azide Free**

<b>Alternate names:</b>	Glucose-6-phosphate dehydrogenase, zwf
<b>Quantity:</b>	10 mg
<b>Concentration:</b>	10.0 mg/ml
<b>Background:</b>	Glucose 6 Phosphate Dehydrogenase (G6PD) produces pentose sugars for nucleic acid synthesis and is the main producer of NADPH reducing power. Catalytic activity: D glucose 6 phosphate + NADP(+) = D glucono 1,5 lactone 6 phosphate + NADPH. Defects in G6PD are the cause of chronic non spherocytic haemolytic anemia (CNSHA). G6PD deficiency is the most common human enzyme deficiency; one benefit of having G6PD deficiency is that it confers a resistance to malaria.
<b>Uniprot ID:</b>	<a href="#">P11412</a>
<b>NCBI:</b>	<a href="#">NP_014158</a>
<b>GeneID:</b>	<a href="#">855480</a>
<b>Host / Isotype:</b>	Rabbit / IgG
<b>Immunogen:</b>	Glucose-6-Phosphate Dehydrogenase isolated and purified from Baker's Yeast. Freund's complete adjuvant is used in the first step of the immunization procedure.
<b>Format:</b>	<b>State:</b> Lyophilized hyperimmune IgG fraction <b>Purification:</b> Ammonium Sulphate Precipitation and Ion Exchange Chromatography <b>Buffer System:</b> PBS, pH 7.2 without preservatives and foreign proteins <b>Reconstitution:</b> Restore by adding 1.0 ml of sterile distilled water
<b>Applications:</b>	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. <i>Recommended Working Dilutions:</i> Non-precipitating antibody-binding techniques: 1/1,000-1/6,000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	This antibody recognizes Glucose-6-Phosphate Dehydrogenase from Baker's Yeast. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. <b>Species:</b> Baker's Yeast. Other species not tested.

**Storage:**

Store lyophilized at 2-8°C for 6 months or at -20°C long term.  
After reconstitution store the antibody undiluted at 2-8°C for one month  
or (in aliquots) at -20°C long term.  
Avoid repeated freezing and thawing.  
Shelf life: one year from despatch.