

**AP21322AF-N****Polyclonal Antibody to beta-Hydroxyacyl-CoA Dehydrogenase - Azide Free**

<b>Alternate names:</b>	HAD, HADHSC, Hydroxyacyl-coenzyme A dehydrogenase mitochondrial, Medium and short chain L-3-hydroxyacyl-coenzyme A dehydrogenase, SCHAD, Short chain 3-hydroxyacyl-CoA dehydrogenase
<b>Quantity:</b>	10 mg
<b>Concentration:</b>	10 mg/ml
<b>Uniprot ID:</b>	<a href="#">P00348</a>
<b>NCBI:</b>	<a href="#">NP_999496</a>
<b>GeneID:</b>	<a href="#">397604</a>
<b>Host / Isotype:</b>	Rabbit / IgG
<b>Immunogen:</b>	beta-Hydroxyacyl-CoA Dehydrogenase isolated and purified from Porcine heart. Freund's complete adjuvant is used in the first step of the immunization procedure.
<b>Format:</b>	<b>State:</b> Lyophilized hyperimmune IgG fraction <b>Purification:</b> Ammonium Sulphate Precipitation and Ion Exchange Chromatography <b>Buffer System:</b> PBS, pH 7.2 without preservatives and foreign proteins <b>Reconstitution:</b> Restore by adding 1.0 ml of sterile distilled water
<b>Applications:</b>	This product is intended for use in precipitating and non-precipitating antibody-binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. <u>Working Dilutions:</u> Non-precipitating antibody-binding techniques: At least 1/100. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
<b>Specificity:</b>	beta-Hydroxyacyl-CoA Dehydrogenase from Porcine heart. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immuno-electrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. <b>Species:</b> Pig. Other species not tested.
<b>Storage:</b>	Store the antibody lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation. Shelf life: one year from despatch.