

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## **OriGene Technologies GmbH**

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## AP21323AF-N Polyclonal Antibody to BDH1 - Azide Free

Alternate names:	3-hydroxybutyrate dehydrogenase, D-beta-hydroxybutyrate dehydrogenase, mitochondrial
Quantity:	10 mg
Concentration:	10 mg/ml
Host / Isotype:	Rabbit / IgG
Immunogen:	beta-Hydroxybutyrate Dehydrogenase isolated and purified from Rhodopseudomonas spheroides. Freund's complete adjuvant is used in the first step of the immunization procedure.
Format:	<b>State:</b> Lyophilized hyperimmune IgG fraction <b>Purification:</b> Ammonium Sulphate Precipitation and Ion Exchange Chromatography <b>Buffer System:</b> PBS, pH 7.2 without preservatives and foreign proteins <b>Reconstitution:</b> Restore by adding 1.0 ml of sterile distilled water
Applications:	This product is intended for use in precipitating and non-precipitating antibody- binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques), to prepare an insoluble immuno-affinity adsorbent, for labelling with a marker of choice. <u>Working Dilutions:</u> Non-precipitating antibody-binding techniques: 1/100-1/500. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	beta-Hydroxybutyrate Dehydrogenase from Rhodopseudomonas spheroides. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. <b>Species:</b> Rhodopseudomonas spheroides. Other species not tested.
Storage:	Store the antibody lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.