

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com OriGene Technologies GmbH

OG/20121030

1/1

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## AP21336BT-N Polyclonal Antibody to LDH1 M4 Isozyme - Biotin

Alternate names:	L-Lactic dehydrogenase LDH-1 Isozyme M4
Quantity:	1 ml
Concentration:	10.0 mg/ml
Host / Isotype:	Rabbit / IgG
Immunogen:	L-Lactic Dehydrogenase LDH-1(M4) isoenzyme, is isolated and purified from Porcine heart. Freund's complete adjuvant is used in the first step of the immunization procedure.
Format:	<ul> <li>State: Lyophilized hyperimmune IgG fraction</li> <li>Purification: Ammonium Sulphate Precipitation and Ion Exchange Chromatography</li> <li>Buffer System: PBS, pH 7.2 without preservatives and foreign proteins</li> <li>Label: Biotin</li> <li>Molar Ratio: Biotin/IgG ~6.4</li> <li>Reconstitution: Restore by adding 1.0 ml of sterile distilled water</li> </ul>
Applications:	This product is intended for use in precipitating and non-precipitating antibody- binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or Histochemical techniques). <u>Working Dilutions:</u> Non-precipitating antibody-binding techniques: 1/1,000-1/7,000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	<ul> <li>L-Lactic Dehydrogenase LDH-1(M4) isoenzyme from Porcine heart.</li> <li>The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and enzyme inhibition.</li> <li>Cross-reactivities against enzymes of other sources may occur but have not been determined.</li> <li>Species: Pig.</li> <li>Other species not tested.</li> </ul>
Storage:	Store the antibody lyophilized at 2-8°C and reconstituted at 2-8°C for one week or (in aliquots) at -20°C for longer. If a slight precipitation occurs upon storage, this should be removed by centrifugation. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.