

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Schillerstr. 5

AP21374BT-N Polyclonal Antibody to Phospholipase A2 - Biotin

Alternate names: Crotoxin basic chain 1, Crotoxin basic chain 2, Phosphatidylcholine 2-acylhydrolase,

Phospholipase A2 CB1, Phospholipase A2 CB2

Quantity: 1 ml

 Concentration:
 10,0 mg/ml

 Uniprot ID:
 P62022

 NCBI:
 8732

Host / Isotype: Rabbit / IgG

Immunogen: Phospholipase A2 isolated and purified from Crotalus durissus terrificus venom.

Freund's adjuvant is used in the first step of the immunization procedure.

Format: State: Lyophilized hyperimmune IgG fraction

Purification: Ammonium Sulphate Precipitation and Ion Exchange Chromatography

Buffer System: PBS, pH 7.2 without preservatives and foreign proteins

Label: Biotin – <u>Marker:</u> N-Hydroxysuccinimidobiotin. <u>Conjugation procedure:</u> A proprietary technique for the binding to biotin is used, followed by several purification steps. After each step activity and specificity are tested in a variety of techniques. The conjugate is lyophilised to assure stability and long shelf life

Molar Ratio: Biotin/IgG ~4.7

Reconstitution: Restore by adding 1.0 ml of sterile distilled water

Applications: Can be used in precipitating and non-precipitating antibody-binding assays (such as

e.g., ELISA and Western blotting and immunofluorescence or histochemical

techniques).

Recommended working dilutions: vary widely, but may be between 1/1000 and

1/20000.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: Biotin-conjugated IgG fraction of polyclonal rabbit antiserum to phospholipase A2

from Crotalus durissus terrificus venom.

The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: immunoelectrophoresis, cross-immunoelectrophoresis, single radial immunodiffusion (Ouchterlony), block titration, ELISA, immunoblotting and

enzyme inhibition.

Storage: Prior to and following reconstitution store the antibody at 2-8°C for one month or at

-20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.