

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

## **OriGene Technologies GmbH**

Schillerstr. 5 32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

## AP21377AF-N Polyclonal Antibody to Phospholipase D - Azide Free

Alternate names:	Choline phosphatase 1, Choline phosphatase 2, PLD-1, PLD-2, PLD1, PLD2, Phosphatidylcholine-hydrolyzing phospholipase D1, Phosphatidylcholine-hydrolyzing phospholipase D2, Phospholipase D alpha 1, Phospholipase D alpha 2
Quantity:	10 mg
Concentration:	10.0 mg/ml
Background:	Phospholipase D (EC=3.1.4.4) is an enzyme involved in lipid degradation by hydrolyzing glycerol phospholipids. Involved in many cellular process including phytohormone action, membrane deterioration, senescence and more.
Uniprot ID:	<u>082549</u>
NCBI:	<u>3716</u>
Host / Isotype:	Rabbit / IgG
Immunogen:	Phospholipase D isolated and purified from Cabbage. Freund's complete adjuvant is used in the first step of the immunization procedure.
Format:	State: Lyophilized hyperimmune IgG fraction Purification: Ammonium Sulphate Precipitation and Ion Exchange Chromatography Buffer System: PBS, pH 7.2 without preservatives and foreign proteins Reconstitution: Restore by adding 1.0 ml of sterile distilled water
Applications:	This product is intended for use in precipitating and non-precipitating antibody- binding assays (such as e.g., ELISA and Western blotting and immunofluorescence or histochemical techniques). To prepare an insoluble Immuno-Affinity adsorbent. For labelling with a marker of the customer's own choice. <u>Working Dilutions:</u> vary widely, but may be between 1/1000 and 1/7000. Other applications not tested. Optimal dilutions are dependent on conditions and should be determined by the user.
Specificity:	IgG fraction of polyclonal Rabbit antiserum to Phospholipase D from Cabbage. The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and Enzyme Inhibition. Cross-reactivities against enzymes of other sources may occur but have not been determined. <b>Species:</b> Cabbage (Brassica oleracea). Other species not tested.
Storage:	Prior to and following reconstitution store the antibody at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing. Shelf life: one year from despatch.

For research and in vitro use only. Not for diagnostic or therapeutic work. Material Safety Datasheets are available at www.acris-antibodies.com or on request.