

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850 UNITED STATES Phone: +1-888-267-4436 Fax: +1-301-340-8606 techsupport@origene.com

OriGene Technologies GmbH

32052 Herford GERMANY Phone: +49-5221-34606-0 Fax: +49-5221-34606-11 info-de@origene.com

Schillerstr. 5

Polyclonal Antibody to Phospholipase D - Aff - Purified AP21377PU-N

Choline phosphatase 1, Choline phosphatase 2, PLD-1, PLD-2, PLD1, PLD2, Alternate names:

Phosphatidylcholine-hydrolyzing phospholipase D1, Phosphatidylcholine-hydrolyzing

phospholipase D2, Phospholipase D alpha 1, Phospholipase D alpha 2

Quantity: 1 mg

Concentration: 2.0 mg/ml

Background: Phospholipase D (EC=3.1.4.4) is an enzyme involved in lipid degradation by

hydrolyzing glycerol phospholipids. Involved in many cellular process including

phytohormone action, membrane deterioration, senescence and more.

Uniprot ID: 082549 NCBI:

Rabbit / IgG **Host / Isotype:**

Phospholipase D isolated and purified from Cabbage. Freund's complete adjuvant is Immunogen:

used in the first step of the immunization procedure.

Format: **State:** Lyophilized purified hyperimmune IgG fraction

Purification: Solid Phase Affinity Chromatography

Buffer System: PBS, pH 7.2 stabilized with Dextran, without preservatives and foreign

prot

3716

Reconstitution: Restore by adding 0.5 ml of sterile distilled water

Applications: This product is intended for use in precipitating and non-precipitating antibody-

> binding assays (such as e.g., ELISA and Western blotting and Immunofluorescence or histochemical techniques). To prepare an insoluble immuno-affinity adsorbent. For

labelling with a marker of the customer's own choice.

Working Dilutions: vary widely, but may be between 1/100 and 1/1000.

Other applications not tested. Optimal dilutions are dependent on conditions and

should be determined by the user.

Specificity: Affinity purified antibodies of polyclonal Rabbit antiserum to Phospholipase D from

Cabbage.

The reagents were evaluated for potency, purity and specificity using most or all of the following techniques: Immunoelectrophoresis, Cross-Immunoelectrophoresis, Single Radial Immunodiffusion (Ouchterlony), block titration, ELISA, Immunoblotting and

Enzyme Inhibition.

Cross-reactivities against enzymes of other sources may occur but have not been

determined.

Species: Cabbage (Brassica oleracea).

Other species not tested.

Storage: Prior to and following reconstitution store the antibody at 2-8°C for one month or at

-20°C for longer.

Avoid repeated freezing and thawing. Shelf life: one year from despatch.